



[REDACTED]

[REDACTED] MSK

[REDACTED] TM

[REDACTED]

[REDACTED] MSK

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED] MSK [REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

MSK ()

MSK 5

:

-
-

SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. The spacer is made of a special material that allows the radiation to pass through it and reach the prostate. The spacer is removed after radiation therapy is completed. This helps to reduce the risk of rectal side effects from radiation therapy.

- SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. The spacer is made of a special material that allows the radiation to pass through it and reach the prostate. The spacer is removed after radiation therapy is completed. This helps to reduce the risk of rectal side effects from radiation therapy. The spacer is made of a special material that allows the radiation to pass through it and reach the prostate. The spacer is removed after radiation therapy is completed. This helps to reduce the risk of rectal side effects from radiation therapy.

SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. The spacer is made of a special material that allows the radiation to pass through it and reach the prostate. The spacer is removed after radiation therapy is completed. This helps to reduce the risk of rectal side effects from radiation therapy. *About Your Fiducial Marker and Rectal Spacer Placement (www.mskcc.org/cancer-care/patient-education/about-placement-fiducial-markers-and-rectal-spacers-radiation-therapy-your-prostate)*

Sperm Banking (www.mskcc.org/cancer-care/patient-education/sperm-banking) *Building Your Family After Cancer Treatment: For People Born With Testicles* (www.mskcc.org/cancer-care/patient-education/building-family-born-with-testicles)

[REDACTED]

[REDACTED]
[REDACTED]
[REDACTED] ([REDACTED])
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED] |

[REDACTED]

[REDACTED]:

- [REDACTED] (Methylcellulose) [REDACTED]
[REDACTED] ([REDACTED] Citrucel® [REDACTED]) |
- [REDACTED] (Simethicone) 125 [REDACTED] ([REDACTED])
[REDACTED] ([REDACTED] Gas-X® Extra Strength) |
- 6 [REDACTED] [REDACTED] ([REDACTED] Fleet® [REDACTED]) |

[REDACTED]
[REDACTED]

[REDACTED]

[REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED] 3 [REDACTED]
[REDACTED]

- [REDACTED] 1 [REDACTED]
(methylcellulose) [REDACTED] | [REDACTED]
[REDACTED]
[REDACTED]

- **Simethicone** (simethicone) **2 (125 mg)** |
- **Simethicone** (simethicone) **2 (125 mg)** |

Simethicone

Simethicone is used to relieve discomfort, such as gas, bloating, and cramping, caused by indigestion, irritable bowel syndrome, or other conditions. It works by breaking up gas bubbles in the stomach and intestines, making it easier to pass. Simethicone is available in various forms, including tablets, capsules, and liquid. It is generally safe and effective when used as directed.

Simethicone is available in various strengths and formulations. The most common strength is 125 mg. It is available in tablets, capsules, and liquid. Simethicone is generally safe and effective when used as directed.

- **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
- **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |

... | ...
...
... 1 ...
... (acetaminophen) (Tylenol®) ...
... |

... |

... |
... |

...

...
... |
... |
... |
... |
... |

... |
... |
... |
... |
... |

... 8 ... 12 ...

- 2 (125 mg) (simethicone) mg

mg

- 1 (methylcellulose) mg
- 2 (125 mg) (simethicone) mg
- 2 mg
- 3 mg
- mg

mg

45 mg 8 mg 12 mg

mg

... ..
... ..
... ..
... ..
... ..

... ..
... .. 15 45
... ..
... ..
... ..
... ..

... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..
... ..
... ..
... ..

- [REDACTED]
- [REDACTED]

[REDACTED]

- [REDACTED] 6 [REDACTED] 8 [REDACTED]
- [REDACTED] 8 [REDACTED]
- [REDACTED]
[REDACTED]:
 - [REDACTED]
 - [REDACTED]
 - [REDACTED]
 - [REDACTED]
- [REDACTED] [REDACTED]

[REDACTED]
[REDACTED], *Improving Your Urinary Continence* (www.mskcc.org/cancer-care/patient-education/improving-your-urinary-continence) [REDACTED]

[REDACTED]
[REDACTED]:

- [REDACTED]

()

-
-
- ()
- (,)
-
-

()

1 2

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED] *Sex and Your Cancer Treatment* (www.mskcc.org/cancer-care/patient-education/sex-cancer-treatment) [REDACTED] | [REDACTED]

[REDACTED]

... | ...
... |
...
...

- ... |
- ... |
..., ...
..., ...
...
- ... |
..., ...
..., ...
..., ...
...

• ... |
...
..., ...
..., ...
...

• ... |
..., ...
..., ...
..., ...
..., ...
... *Managing Cancer-Related Fatigue with Exercise* (www.mskcc.org/cancer-care/patient-education/managing-cancer-related-fatigue-exercise) ...
..., ...
..., ...
...

- ...

- [REDACTED]
- [REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]?

[REDACTED]?

[REDACTED]

