



[REDACTED]

[REDACTED] MSK

[REDACTED] TM

[REDACTED]

[REDACTED] MSK

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED] MSK [REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

MSK ()

MSK 5

:

-
-

SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum.

- SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum. (SpaceOAR® is a registered trademark of SpaceOAR, Inc.) SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum. 3 SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum. (SpaceOAR® is a registered trademark of SpaceOAR, Inc.) SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum.

SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum. *About Your Fiducial Marker and Rectal Spacer Placement* (www.mskcc.org/cancer-care/patient-education/about-placement-fiducial-markers-and-rectal-spacers-radiation-therapy-your-prostate)

SpaceOAR® is a non-invasive, temporary spacer that is placed between the prostate and the rectum before radiation therapy. It is made of a special material that is safe and effective. It is used to reduce the risk of side effects from radiation therapy to the rectum. *Sperm Banking* (www.mskcc.org/cancer-care/patient-education/sperm-banking) *Building Your Family After Cancer Treatment: For People Born With Testicles* (www.mskcc.org/cancer-care/patient-education/building-family-born-with-testicles)

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX

XX
XX
XX (XX)
XX
XX |
XX
XX
XX
XX
XX
XX
XX
XX
XX
XX |

XXXXXXXXXX

XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX:

- XXXXXXXXXXXXXXXXXXXXXXXXXXXXX (Methylcellulose) XXXXXXXXXXXXXXXXXXXX
 XXXXXXXXXXXXXXXXXXXX (XXXXXX Citrucel® XXXXXXXXXXXX) |
- XXXXXXXXXXXXXXXXXXXXX (Simethicone) 125 XXXXXXXXXXXXXXXXXXXX (XXXXXXXXXX)
 XXXXXXXXXXXXXXXXXXXX (XXXXXX Gas-X® Extra Strength) |
- 6 XXXXXXXXXXXXXXXXXXXXXXXXXXXX (XXXXXX Fleet® XXXXXXXXXXXXXXXXXXXX
 XXXXXXXXXXXX) |

XX
XX |

XXXXXXXXXXXXXXXXXXXXXXXXXXXX

XXXX XXXX XXXXXXXXXXX XXXX XX
XXXX XXX
XX |
XX 3 XXXXXXXXXXX
XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX |

- XXXXXXXXXXXXXXX 1 XXX
 (methylcellulose) XXXXXXXXXXXXXXXXXXXX | XXXXXXXXXXXXXXXXXXXX
 XXX
 XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX |

- **Simethicone** (simethicone) **2 (125 mg)** |
- **Simethicone** (simethicone) **2 (125 mg)** |

Simethicone (simethicone) **2 (125 mg)** |

Simethicone (simethicone) 2 (125 mg) is used to relieve gas and bloating. It works by breaking up gas bubbles in the stomach and intestines. It is available as a liquid suspension and as tablets. It is generally safe and effective. Common side effects include nausea and vomiting. It should be used with caution in children and pregnant women. It is not recommended for use in patients with severe liver disease. It is also not recommended for use in patients with severe kidney disease. It is important to follow the instructions on the label and to consult with a healthcare provider if you have any questions or concerns.

Simethicone (simethicone) 2 (125 mg) is used to relieve gas and bloating. It works by breaking up gas bubbles in the stomach and intestines. It is available as a liquid suspension and as tablets. It is generally safe and effective. Common side effects include nausea and vomiting. It should be used with caution in children and pregnant women. It is not recommended for use in patients with severe liver disease. It is also not recommended for use in patients with severe kidney disease. It is important to follow the instructions on the label and to consult with a healthcare provider if you have any questions or concerns.

- **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
- **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |
 - **Simethicone** (simethicone) **2 (125 mg)** |

[Redacted text block]

[Redacted text block]

(RDA)- [Redacted text]

[Redacted text block]

[Redacted text block]

- [Redacted text] (CGM)
- [Redacted text]

[Redacted text block]

[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED] 1 [REDACTED]
[REDACTED] (acetaminophen) (Tylenol®) [REDACTED]
[REDACTED]

[REDACTED], [REDACTED]
[REDACTED]

[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED] | [REDACTED], [REDACTED]
[REDACTED]

[REDACTED]

[REDACTED]
[REDACTED], [REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED] [REDACTED] [REDACTED] [REDACTED] [REDACTED]
[REDACTED] [REDACTED] | [REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]

[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]

[REDACTED]
[REDACTED] 8 [REDACTED] 12 [REDACTED]

- 2 (125 mg) (simethicone) mg

mg

- 1 (methylcellulose) mg
- 2 (125 mg) (simethicone) mg
- 2 mg
- 3 mg
- mg

mg

45 mg 8 mg 12 mg

mg

- 6 weeks after surgery, you may be able to return to normal activities.
- You may experience some of the following side effects:

Weakness, fatigue, and weight loss are common after surgery. You may also have changes in your appearance, such as a lump or swelling in the groin area. It's important to take care of yourself and get plenty of rest. You may also have some changes in your diet. You may be able to eat a normal diet soon after surgery, but you may need to eat a soft diet for a few days. You may also have some changes in your bowel habits. You may have some constipation or diarrhea. You may also have some changes in your urinary habits. You may have some incontinence or urgency. You may also have some changes in your sexual function. You may have some problems with erection or orgasm.

- You may have some of the following side effects:
 - Weakness, fatigue, and weight loss
 - Changes in your appearance, such as a lump or swelling in the groin area
 - Changes in your diet
 - Changes in your bowel habits
 - Changes in your urinary habits
 - Changes in your sexual function
- You may also have some of the following side effects:
 - Constipation
 - Diarrhea
 - Incontinence
 - Urgency
- You may also have some of the following side effects:
 - Problems with erection or orgasm

For more information on these side effects, please see our patient education material, *Improving Your Urinary Continence* (www.mskcc.org/cancer-care/patient-education/improving-your-urinary-continence)

You may also have some of the following side effects:

- Incontinence
- Urgency

- You may also have some of the following side effects:
 - Incontinence
 - Urgency

... | ...
... |
...
...

- ...

- ...

- ...

- ...

- ...
... *Managing Cancer-Related Fatigue with Exercise* (www.mskcc.org/cancer-care/patient-education/managing-cancer-related-fatigue-exercise) ...

- ...

1. 本公司 (AAA - PSA 有限公司), 係一家
 在 [] 年 [] 月 [] 日 注册成立的有限公司, 注册
 资本为 [] 元, 注册地址为 [] 省 [] 市 [] 区 [] 路 [] 号。
 2. 本公司经营的主要业务为 [] 产品的研发、生产、销售。
 3. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。

(根据 [] 年 [] 月 [] 日 签订的 [] 合同) 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 4. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。

5. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。

- 1. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 2. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 3. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 4. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 5. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
- 6. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 7. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 8. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 9. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。
 10. 根据 [] 年 [] 月 [] 日 签订的 [] 合同, 本公司
 向 [] 公司 销售了 [] 台 [] 产品, 总金额为 [] 元。

6. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。
 7. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。
 8. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。

9. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。
 10. 本公司在 [] 年 [] 月 [] 日 至 [] 年 [] 月 [] 日 期间, 共
 销售了 [] 台 [] 产品, 总金额为 [] 元。

[REDACTED]:

- [REDACTED]
- [REDACTED]
- [REDACTED]

[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]

[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED] | [REDACTED]
[REDACTED]:

- [REDACTED]
[REDACTED]
- [REDACTED]
[REDACTED]
[REDACTED]
- [REDACTED]
[REDACTED]

[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]:

- [REDACTED]
[REDACTED]

(ED) [REDACTED]

- [REDACTED]
- [REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]

[REDACTED]?

[REDACTED]?

[REDACTED]

	<p>■■■■■■■■■■■■■■■■■■■■■ </p>	
<p>■■■■■■■■■■■■■■■■■■■■■</p>	<ul style="list-style-type: none"> ● ■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■ ■■■■■■■■■■ 2 ■■■■■■ 3 ■■■■■■■■■■ ■■■■■■■■■■ ■■■■■■ ■■■■■■■■■■■■ ■■■■■■■■■■ ■■■■■■ 	
<p>■■■■■■■</p>	<ul style="list-style-type: none"> ● ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■ ■■■■■■ ■■■■■■■■■■ ■■■■■■■■■■■■■■■■ ■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■ ■■■■■■ 	
<p>■■■■■</p>	<ul style="list-style-type: none"> ● ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■ ■■■■■■ ● ■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■ 1 ■■■■■■■■ ■■■■ ■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■ (acetaminophen) (Tylenol®) ■■■ ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■ 	<p>■■■■■■■■■■■■■ ■■■■■■■■■■■■ ■■■■■■■■■■ ■■■■■ </p>

■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■

Please arrive at the hospital at least 15 minutes before your appointment. If you are late, your appointment may be canceled. You will be given a form to fill out at the hospital. Please bring a list of all the medications you are taking, including over-the-counter medications, vitamins, and herbal supplements. You will also need to bring a list of all the symptoms you are experiencing and a list of all the tests you have had in the past year.

9:00 AM - 5:00 PM

Patient name: _____
 Phone number: _____

Patient ID: _____
 Date of birth: _____

MSK Precise® Radiation Therapy for Prostate Cancer is a new, highly personalized treatment. It uses a combination of external beam radiation therapy and brachytherapy to deliver a precise dose of radiation to the prostate gland. This treatment is designed to be more effective than traditional radiation therapy and has fewer side effects.

MSK Precise® Radiation Therapy for Prostate Cancer is available at Memorial Sloan Kettering Cancer Center. For more information, please contact our Radiation Therapy Department at 212-639-2000.

For more information, please visit our website at www.mskcc.org/pe.

About Your MSK Precise® Radiation Therapy for Prostate Cancer - Last updated on June 13, 2024
 Memorial Sloan Kettering Cancer Center