

PATIENT & CAREGIVER EDUCATION

Zinc Oxide

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

AmeriDerm PeriShield [OTC]; Ammens Original Medicated [OTC]; Ammens Shower Fresh [OTC]; Balmex [OTC]; Baza Protect Moisture Barrier [OTC]; Boudreauxs Butt Paste [OTC]; Calmol 4 [OTC]; CareAll [OTC]; Critic-Aid Thick Moist Barrier [OTC]; Desitin Maximum Strength Original [OTC]; Desitin Rapid Relief [OTC]; Dr. Smith's Adult Barrier [OTC] [DSC]; Dr. Smith's Diaper [OTC]; Endit [OTC]; GoodSense Diaper Rash [OTC]; Medpura [OTC]; PeriGuard [OTC] [DSC]; Pharmabase Barrier [OTC]; Secura Extra Protective [OTC]; Secura Protective [OTC]; Triple Paste [OTC]; Z-Guard [OTC]

Brand Names: Canada

Zincofax®

What is this drug used for?

All skin products:

- It is used to treat or prevent diaper rash.
- It is used to treat skin irritation.
- It is used to treat poison oak, ivy, and sumac rashes.

Suppository:

- It is used to treat signs of hemorrhoids or rectal irritation.

What do I need to tell the doctor BEFORE my child takes this drug?

All products:

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

All skin products:

- If your child has a deep wound or a puncture wound.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not use more often or for longer than you were told.
- This drug may stain clothing or fabric. Protect clothing and fabrics from staining.
- This drug may cause harm if swallowed. If this drug is swallowed, call a doctor or poison control center right away.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if your child has any side effects that bother your child or do not go away.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All skin products:

- Do not give by mouth. Use on your child's skin only. Keep out of your child's mouth, nose, and eyes (may burn).
- Wash your hands before and after use.
- Clean affected part before use. Make sure to dry well.
- Put a thin layer on the affected part and rub in gently.
- If you are using under a diaper, change the diaper often. Clean the affected area and allow it to dry. Use the drug with each diaper changed, especially at bedtime.

Suppository:

- This drug is for rectal use only. Do not give this drug by mouth.
- Wash your hands before and after use.
- If possible, clean the area before using this drug. Pat or blot with a cleansing pad, then pat dry with toilet tissue or a soft cloth. Ask your doctor if you are not sure what kind of cleansing pad to use.
- Take foil off the rectal suppository and put in, pointed end first. Do not handle too much.
- If suppository is soft, chill in a refrigerator or run cold water over it.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- Many times this drug is given on an as needed basis. Do not give to your child more often than told by the doctor.

How do I store and/or throw out this drug?

All products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Suppository:

- Protect from heat.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical

advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-06-03

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Zinc Oxide - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center