



PATIENT & CAREGIVER EDUCATION

About Your Colposcopy

This information describes what to expect during and after your colposcopy.

A colposcopy is a procedure that examines your cervix, which is located at the end of your uterus. This procedure is done to get a close-up look of your cervix to check for cancer..




Figure 1. Female anatomy

During your colposcopy, your healthcare provider will use a tool called a colposcope. This tool is like a magnifying glass for your cervix and will help your healthcare provider find any signs of cancer.

Your healthcare provider may also do a biopsy. This means they will take a small sample of tissue from your cervix. You may also have another type of biopsy done called an endocervical curettage (ECC). During an ECC, your healthcare provider will remove cells from the lining of your endocervical canal (canal that goes from your cervix to your uterus).

Your healthcare provider will send the samples to a lab to be checked for cancer.

The procedure usually takes about 20 minutes.

Before your procedure

Schedule your procedure for 1 week after your period.

The day before your procedure, do not:

- Have vaginal intercourse.
- Put anything in your vagina, including douches and tampons.
- Use any vaginal creams or medications.

You may eat before your procedure.

During your procedure

- You will lie on your back with your feet in stirrups. This is the same position you're in for your routine pelvic exams.
- Your healthcare provider will put a speculum into your vagina. This is an instrument that will gently spread apart your vaginal walls so that your healthcare provider can see your cervix.
- Your healthcare provider will wash your cervix with a mild vinegar solution. This will clear any mucus on your cervix and make it easier to see any abnormal cells.
- Then your healthcare provider will then examine your cervix using the colposcope.
- If your healthcare provider performs a biopsy, you will feel some discomfort as the tissue is removed.
- If your healthcare provider performs an ECC, you will feel mild cramping.

After your procedure

If you had a biopsy or ECC:

- You may have a bloody or brownish-black vaginal discharge for a few days. Use sanitary pads for vaginal discharge. Do not use a tampon.
- Take acetaminophen (Tylenol®) or ibuprofen (Advil®) for any discomfort.
- For 1 week after your procedure, or as instructed by your doctor, do not:
 - Have vaginal intercourse.
 - Put anything in your vagina, including douches and tampons.
 - Use any vaginal creams or medications.
- Call your healthcare provider's office 1 to 2 weeks after your procedure for your biopsy results.

Call your healthcare provider if you have:

- A fever of 101° F (38.3° C) or higher.
- Vaginal discharge lasting more than a few days.
- Vaginal bleeding that is heavier than your menstrual flow.
- Pain that doesn't get better with over the counter pain medication.
- Any unexpected problems.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

About Your Colposcopy - Last updated on September 6, 2024
All rights owned and reserved by Memorial Sloan Kettering
Cancer Center