



About Your Ileostomy Closure Surgery

This guide will help you get ready for your ileostomy closure surgery at MSK. It will also help you know what to expect as you recover.

Use this guide as a source of information in the days leading up to your surgery. Bring it with you on the day of your surgery. You and your care team will use it as you learn more about your recovery.

Your surgery

Surgery date: _____

Surgery location: Presurgical Center (PSC) on the 6th floor
1275 York Ave. (between East 67th and East 68th streets)
New York, NY 10065
Take the B elevator to the 6th floor.

Expected discharge date: _____

It's best to plan your ride home from the hospital ahead of time. This will help you keep from waiting for your ride after you're discharged from (leave) the hospital. For more information, read the section "Leaving the hospital."



Visit www.msk.org/pe/ileostomy-closure to view this guide online.

Table of contents

About your ileostomy closure surgery	3
About your ileostomy	3
About your ileostomy closure surgery	3
Before your ileostomy closure surgery	5
Getting ready for your surgery	6
Within 30 days of your surgery	10
7 days before your surgery	12
2 days before your surgery	13
1 day before your surgery	14
The day of your surgery	16
After your ileostomy closure surgery	21
In the Post-Anesthesia Care Unit (PACU)	22
In your hospital room	23
At home	26
When to call your healthcare provider	35
Support services	37
MSK support services	38
External support services	43
Educational resources	45
<i>Herbal Remedies and Cancer Treatment</i>	
<i>How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil</i>	

About your ileostomy closure surgery

About your ileostomy

An ileostomy is the opening your surgeon made in your abdomen (belly) when you had surgery to remove part of your colon or rectum. Your ileostomy was created from the part of your small intestine called the ileum. It let your bowel movements (poop) pass out of your body and into a stoma bag. Your ileostomy was made so your anastomosis (the place where your surgeon reconnected your bowel) could heal.

About your ileostomy closure surgery

An ileostomy closure surgery is done to reverse your ileostomy so you can have bowel movements like you did before your surgery. Ileostomy closure surgery is usually done through your stoma (see Figure 1). Your surgeon may need to make an additional incision (surgical cut), but this is rare.

After your surgery, you'll have a small wound where your ileostomy used to be. This wound will heal in about 4 to 6 weeks. You'll need to change the bandage on this wound every day. While you're in the hospital, your nurse will teach you how to change your bandage and give you the supplies you'll need to do it at home.




Figure 1. Your stoma

Notes _____

Before your ileostomy closure surgery

This section will help you get ready for your surgery. Read it when your surgery is scheduled. Refer to it as your surgery gets closer. It has important information about what to do to get ready.



As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

Getting ready for your surgery

You and your care team will work together to get ready for your surgery. Help us keep you safe by telling us if any of these things apply to you, even if you're not sure.

- I take prescription any medicines. A prescription medicine is one you can only get with a prescription from your healthcare provider. Examples include:
 - Medicines you swallow.
 - Medicines you take as an injection (shot).
 - Medicines you inhale (breathe in).
 - Medicines you put on your skin as a patch or cream.
- I take over-the-counter medicines, including patches and creams. An over-the-counter medicine is one you can buy without a prescription.
- I take dietary supplements, such as herbs, vitamins, minerals, or natural or home remedies.
- I have a pacemaker, automatic implantable cardioverter-defibrillator (AICD), or other heart device.
- I have had a problem with anesthesia (A-nes-THEE-zhuh) in the past. Anesthesia is medicine to make you sleep during a surgery or procedure.
- I'm allergic to certain medicines or materials, including latex.
- I'm not willing to receive a blood transfusion.
- I use recreational drugs, such as marijuana.

Always be sure your healthcare providers know all the medicines and supplements you're taking.

You may need to follow special instructions before surgery based on the medicines and supplements you take. If you do not follow those instructions, your surgery may be delayed or canceled.

About drinking alcohol

It's important to talk with your healthcare providers about how much alcohol you drink. This will help us plan your care.

If you drink alcohol regularly, you may be at risk for problems during and after your surgery. These include bleeding, infections, heart problems, and a longer hospital stay.

If you drink alcohol regularly and stop suddenly, it can cause seizures, delirium, and death. If we know you're at risk for these problems, we can prescribe medicine to help prevent them.

Here are things you can do before surgery to keep from having problems.

- Be honest with us about how much alcohol you drink.
- Try to stop drinking alcohol once your surgery is planned. Tell us right away if you:
 - Get a headache.
 - Feel nauseous (like you're going to throw up).
 - Feel more anxious (nervous or worried) than usual.
 - Cannot sleep.

These are early signs of alcohol withdrawal and can be treated.

- Tell us if you cannot stop drinking.
- Ask us questions about drinking and surgery. We will keep all your medical information private, as always.

About smoking

If you smoke or use an electronic smoking device, you can have breathing problems when you have surgery. Vapes and e-cigarettes are examples of electronic smoking devices. Stopping for even a few days before surgery can help prevent breathing problems during and after surgery.

We will refer you to our Tobacco Treatment Program if you smoke. You can also reach the program by calling 212-610-0507. To learn more, visit www.msk.org/tobacco

About sleep apnea

Sleep apnea is a common breathing problem. If you have sleep apnea, you stop breathing for short lengths of time while you're asleep. The most common type is obstructive sleep apnea (OSA). With OSA, your airway becomes fully blocked during sleep.

OSA can cause serious problems during and after surgery. Tell us if you have or think you might have sleep apnea. If you use a breathing device, such as a CPAP machine, bring it on the day of your surgery.

Using MSK MyChart

MSK MyChart (mskmychart.mskcc.org) is MSK's patient portal. You can use it to send and read messages from your care team, view your test results, see your appointment dates and times, and more. You can also invite your caregiver to make their own account so they can see information about your care.

If you do not have an MSK MyChart account, you can sign up at mskmychart.mskcc.org. You can also ask a member of your care team to send you an invitation.

If you need help with your account, call the MSK MyChart Help Desk at 646-227-2593. They are available Monday through Friday between 9 a.m. and 5 p.m. (Eastern time).

About your Goals to Discharge Checklist

After your surgery, you'll focus on getting well enough to leave the hospital. We'll send a Goals to Discharge Checklist to your MSK MyChart account to help you track how you're doing.

You can use your Goals to Discharge Checklist to see the goals you need to meet before leaving the hospital. You can also update your progress throughout the day. Your updates send alerts to your surgical team about your progress.

To learn more, read *Frequently Asked Questions About the Goals to Discharge Checklist*. You can ask for a printed copy or find it at www.msk.org/pe/goals_discharge_checklist

About Enhanced Recovery After Surgery (ERAS)

ERAS is a program to help you get better faster after your surgery. It's important to do certain things before and after your surgery as part of the ERAS program.

Before your surgery, make sure you're ready by:

- **Reading this guide.** It will help you know what to expect before, during, and after your surgery. If you have questions, write them down. You can ask your healthcare provider at your next visit or call their office.
- **Exercising and following a healthy diet.** This will help get your body ready for your surgery.

After your surgery, help yourself recover more quickly by:

- **Reading your recovery pathway.** This is an educational resource your healthcare provider will give you. It has goals for your recovery. It will help you know what to do and expect each day.
- **Starting to move around as soon as you can.** The sooner you get out of bed and walk, the quicker you can get back to your usual activities.

Within 30 days of your surgery

Presurgical testing (PST)

You'll have a PST appointment before your surgery. You'll get a reminder from your surgeon's office with the appointment date, time, and location. Visit www.msk.org/parking for parking information and directions to all MSK locations.

You can eat and take your usual medicine(s) the day of your appointment.

It's helpful to bring these things to your appointment:

- A list of all the medicines you're taking, including prescription and over-the-counter medicines, patches, and creams.
- Results of any medical tests done outside of MSK in the past year, if you have them. Examples include results from a cardiac stress test, echocardiogram, or carotid doppler study.
- The names and telephone numbers of your healthcare providers.

You'll meet with an advance practice provider (APP) during your PST appointment. They work closely with MSK's anesthesiology (A-nes-THEE-zee-AH-loh-jee) staff. These are healthcare providers with special training in using anesthesia during a surgery or procedure.

Your APP will review your medical and surgical history with you. You may have tests to plan your care, such as:

- An electrocardiogram (EKG) to check your heart rhythm.
- A chest X-ray.
- Blood tests.

Your APP may recommend you see other healthcare providers. They'll also talk with you about which medicine(s) to take the morning of your surgery.

Identify your caregiver

Your caregiver has an important role in your care. Before your surgery, you and your caregiver will learn about your surgery from your healthcare providers. After your surgery, your caregiver will take you home when you're discharged. They'll also help you care for yourself at home.



For caregivers

Caring for a person going through cancer treatment comes with many responsibilities. We offer resources and support to help you manage them.

Visit www.msk.org/caregivers or read *A Guide for Caregivers* to learn more. You can ask for a printed copy or find it at www.msk.org/pe/guide_caregivers

Fill out a Health Care Proxy form

If you have not already filled out a Health Care Proxy form, we recommend you do now. If you already filled one out or have any other advance directives, bring them to your next appointment.

A health care proxy is a legal document. It says who will speak for you if you can't communicate for yourself. This person is called your health care agent.

- To learn about health care proxies and other advance directives, read *Advance Care Planning for People With Cancer and Their Loved Ones*. You can find it at www.msk.org/pe/advance_care_planning or ask for a printed copy.
- To learn about being a health care agent, read *How to Be a Health Care Agent*. You can find it at www.msk.org/pe/health_care_agent or ask for a printed copy.

Talk with a member of your care team if you have questions about filling out a Health Care Proxy form.

Buy clear liquids

You'll need to follow a clear liquid diet before your surgery. It's helpful to buy your supplies ahead of time. For a list of clear liquids you can drink, read the section "Follow a clear liquid diet."

Buy acetaminophen (Tylenol®) and ibuprofen (Advil® or Motrin®)

Acetaminophen and ibuprofen are over-the-counter pain medicines. You'll use them after your surgery to help manage your pain at home.

It's helpful to buy these medicines ahead of time if you don't already have them. You can get them at your local pharmacy without a prescription.

7 days before your surgery

Follow your healthcare provider's instructions for taking aspirin

Aspirin can cause bleeding. If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless they tell you to.**

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the "Educational resources" section of this guide.

Stop taking vitamin E, multivitamins, herbal remedies, and other dietary supplements

Vitamin E, multivitamins, herbal remedies, and other dietary supplements can cause bleeding. Stop taking them 7 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *Herbal Remedies and Cancer Treatment*. You can find it in the “Educational resources” section of this guide.

2 days before your surgery

Stop taking nonsteroidal anti-inflammatory drugs (NSAIDs)

NSAIDs, such as ibuprofen (Advil® and Motrin®) and naproxen (Aleve®), can cause bleeding. Stop taking them 2 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the “Educational resources” section of this guide.

1 day before your surgery

Follow a clear liquid diet

You'll need to follow a clear liquid diet the day before your surgery. A clear liquid diet includes only liquids you can see through. You can find examples in the "Clear liquid diet" table.

While you're following a clear liquid diet:

- Do not eat any solid foods.
- Try to drink at least 1 (8-ounce) cup of clear liquid every hour you're awake.
- Drink different types of clear liquids. Do not just drink water, coffee, and tea.
- Do not drink any liquids you can't see through, such as milk or smoothies.
- Do not drink sugar-free liquids unless you have diabetes and a member of your care team tells you to.

How to follow a clear liquid diet if you have diabetes

Ask the healthcare provider who manages your diabetes:

- What to do while you're following a clear liquid diet.
- If you need to change your dose of insulin or other diabetes medicine(s), if you take them.
- If you should drink sugar-free clear liquids.

Check your blood sugar level often while you're following a clear liquid diet. If you have questions, talk with your healthcare provider.

Clear liquid diet

	OK to have	Do not have
Soups	<ul style="list-style-type: none"> • Clear broth, bouillon, and consommé. 	<ul style="list-style-type: none"> • Anything with pieces of food or seasoning.
Sweets	<ul style="list-style-type: none"> • Gelatin, such as Jell-O®. • Flavored ices. • Hard candies, such as Life Savers®, lemon drops, and peppermints. 	<ul style="list-style-type: none"> • All other sweets.
Drinks	<ul style="list-style-type: none"> • Clear fruit juices, such as lemonade, apple, cranberry, and grape juices. • Soda, such as ginger ale, 7UP®, Sprite®, and seltzer. • Sports drinks, such as Gatorade® and Powerade®. • Black coffee or plain tea without any type of milk or creamer. • Water, including carbonated (fizzy) and flavored water. • Clear nutritional drinks, such as Boost® Breeze, Ensure Clear™, Pedialyte®, and Diabetishield®. 	<ul style="list-style-type: none"> • Juices with pulp. • Nectars. • Smoothies or shakes. • Milk, cream, and other dairy products. • Nut milks, plant milks, non-dairy creamers, and other dairy alternatives. • Drinks with alcohol.

Note the time of your surgery

A staff member will call you after 2 p.m. the day before your surgery. If your surgery is scheduled for a Monday, they'll call you the Friday before. If you do not get a call by 7 p.m., call 212-639-5014.

The staff member will tell you what time to get to the hospital for your surgery. They'll also remind you where to go.

Visit www.msk.org/parking for parking information and directions to all MSK locations.

Instructions for eating



Stop eating at midnight (12 a.m.) the night before your surgery. This includes hard candy and gum.

Your healthcare provider may have given you different instructions for when to stop eating. If so, follow their instructions. Some people need to fast (not eat) for longer before their surgery.

The day of your surgery

Instructions for drinking

Between midnight (12 a.m.) and 2 hours before your arrival time, only drink the liquids on the list below. Do not eat or drink anything else. Stop drinking 2 hours before your arrival time.

- Water.
- Clear apple juice, clear grape juice, or clear cranberry juice.
- Gatorade or Powerade.
- Black coffee or plain tea. It's OK to add sugar. Do not add anything else.
 - Do not add any amount of any type of milk or creamer. This includes plant-based milks and creamers.
 - Do not add honey.

- Do not add flavored syrup.

If you have diabetes, pay attention to the amount of sugar in your drinks. It will be easier to control your blood sugar levels if you include sugar-free, low-sugar, or no added sugar versions of these drinks.

It's helpful to stay hydrated before surgery, so drink if you are thirsty. Do not drink more than you need. You will get intravenous (IV) fluids during your surgery.



Stop drinking 2 hours before your arrival time. This includes water.

Your healthcare provider may have given you different instructions for when to stop drinking. If so, follow their instructions.

Take your medicines as instructed

A member of your care team will tell you which medicines to take the morning of your surgery. Take only those medicines with a sip of water. Depending on what you usually take, this may be all, some, or none of your usual morning medicines.

Things to remember

- Wear something comfortable and loose-fitting.
- If you wear contact lenses, wear your glasses instead. Wearing contact lenses during surgery can damage your eyes.
- Do not wear any metal objects. Take off all jewelry, including body piercings. The tools used during your surgery can cause burns if they touch metal.
- Leave valuable items at home.

- If you're menstruating (have your monthly period), use a sanitary pad, not a tampon. We'll give you disposable underwear and a pad if you need them.

What to bring

- A pair of loose-fitting pants (such as sweatpants).
- Sneakers that lace up. You may have some swelling in your feet. Lace-up sneakers can fit over this swelling.
- Your breathing device for sleep apnea (such as your CPAP machine), if you have one.
- Your Health Care Proxy form and other advance directives, if you filled them out.
- Your cell phone and charger.
- Only the money you may want for small purchases, such as a newspaper.
- A case for your personal items, if you have any. Eyeglasses, hearing aids, dentures, prosthetic devices, wigs, and religious articles are examples of personal items.
- This guide. You'll use it to learn how to care for yourself after surgery.

Once you're in the hospital

When you get to the hospital, take the B elevator to the 6th floor. Check in at the desk in the PSC waiting room.

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having surgery on the same day.

We'll give you a hospital gown, robe, and nonskid socks to wear when it's time to change for surgery.

Meet with a nurse

You'll meet with a nurse before surgery. Tell them the dose of any medicines you took after midnight (12 a.m.) and the time you took them. Make sure to include prescription and over-the-counter medicines, patches, and creams.

Your nurse may place an intravenous (IV) line in one of your veins, usually in your arm or hand. If your nurse does not place the IV, your anesthesiologist (A-nes-THEE-zee-AH-loh-jist) will do it in the operating room.

Meet with an anesthesiologist

You'll also meet with an anesthesiologist before surgery. They will:

- Review your medical history with you.
- Ask if you've had any problems with anesthesia in the past, such as nausea or pain.
- Talk with you about your comfort and safety during your surgery.
- Talk with you about the kind of anesthesia you'll get.
- Answer your questions about your anesthesia.

Get ready for surgery

When it's time for your surgery, you'll take off your eyeglasses, hearing aids, dentures, prosthetic devices, wig, and religious articles.

You'll either walk into the operating room or a staff member will bring you there on a stretcher. A member of the operating room team will help you onto the operating bed. They'll put compression boots on your lower legs. These gently inflate and deflate to help blood flow in your legs.

Once you're comfortable, your anesthesiologist will give you anesthesia through your IV line and you'll fall asleep. You'll also get fluids through your IV line during and after your surgery.

During your surgery

After you're fully asleep, your care team will place a breathing tube through your mouth into your airway. Your breathing tube is usually taken out while you're still in the operating room.

Notes _____

After your ileostomy closure surgery

This section will help you know what to expect after your surgery. You'll learn how to safely recover from your surgery both in the hospital and at home.



As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

In the Post-Anesthesia Care Unit (PACU)

You'll be in the PACU when you wake up after your surgery. A nurse will be keeping track of your temperature, pulse, blood pressure, and oxygen levels. You may get oxygen through a tube resting below your nose or a mask over your nose and mouth. You'll also have compression boots on your lower legs.

Pain medicine

You'll get medicine to control your pain and keep you comfortable. There are different ways pain medicine can be given:

- **Nerve block:** Some people get a nerve block before or during surgery. With a nerve block, your healthcare provider injects medicine into some of your nerves to reduce pain after surgery.
- **IV medicine:** Some people get pain medicine into a vein through their IV line.

You'll have 1 or more of these after your surgery. They're all effective methods to control your pain. Your healthcare provider will talk with you before choosing the best one(s) for you.

Moving to your hospital room

You'll stay in the PACU until you're awake and your pain is under control. Most people move to their hospital room after a few hours in the PACU, but some people stay in the PACU overnight for observation.

After your stay in the PACU, a staff member will take you to your hospital room.

In your hospital room

The length of time you're in the hospital after your surgery depends on your recovery. Most people stay in the hospital for 2 days. Your care team will tell you what to expect.

In your hospital room, you'll meet one of the nurses who will care for you during your stay. A nurse will help you out of bed and into your chair soon after you get there.

Your care team will teach you how to care for yourself while you're healing from your surgery. You can help yourself recover more quickly by:

- **Reading your recovery pathway.** We will give you a pathway with goals for your recovery if you do not already have one. It will help you know what to do and expect on each day during your recovery.
- **Starting to move around as soon as you can.** The sooner you get out of bed and walk, the quicker you can get back to your usual activities.

You can use your Goals to Discharge Checklist to track your progress during your recovery. Read *Frequently Asked Questions About the Goals to Discharge Checklist* to learn more. You ask for a printed copy or find it at www.msk.org/pe/goals_discharge_checklist

Managing your pain

You'll have some pain after your surgery. Your healthcare providers will ask you about your pain often and give you medicine as needed. If your pain isn't relieved, tell one of your healthcare providers. It's important to control your pain so you can use your incentive spirometer and move around. Controlling your pain will help you recover better.

Many people find their pain is controlled with over-the-counter medicines alone. If you need stronger pain medicine in the hospital, one of your healthcare providers will give you a prescription before you leave. Talk with your healthcare providers about possible side effects and how to taper (slowly stop taking) your medicine.

Moving around and walking

Moving around and walking will help lower your risk for blood clots and pneumonia (lung infection). It will also help you start passing gas and having bowel movements (pooping) again.

Read your recovery pathway to learn about your specific moving and walking goals. Your nurse, physical therapist, or occupational therapist will help you move around, if needed.

To learn more about how walking can help you recover, read *Frequently Asked Questions About Walking After Your Surgery*. You can find it at www.msk.org/pe/walking_after_surgery or ask for a printed copy.

Read *Call! Don't Fall!* to learn what you can do to stay safe and keep from falling while you're in the hospital. You can ask for a printed copy or find it at www.msk.org/pe/call_dont_fall

Exercising your lungs

It's important to exercise your lungs so they expand fully. This helps prevent pneumonia.

- Your nurse will give you an incentive spirometer. Use it 10 times every hour you're awake. Read *How To Use Your Incentive Spirometer* to learn more. You can find it at www.msk.org/pe/incentive_spirometer or ask for a printed copy.

- Do coughing and deep breathing exercises. A member of your care team will teach you how.

Eating and drinking

Right after your surgery, you will not be able to eat solid foods. You'll follow a liquid diet. After that, you'll slowly start eating solid foods. This usually happens the day after your surgery.

A clinical dietitian nutritionist will give you more information about eating after surgery before you leave the hospital.

Caring for your wound

You'll have a wound where your ileostomy used to be. It will not be closed with sutures (stitches) or staples, and it will be very raw and red. This is normal.

Your nurse will pack (fill) your wound with gauze and change your bandage every day while you're in the hospital. They'll also teach you how to do it yourself. They will have you practice packing your wound and changing your dressing before you go home.

It's helpful if your caregiver also learns how to care for your wound. This will make it easier for them to help you care for yourself at home.

Your wound should close 4 to 6 weeks after your surgery.

Planning for discharge

Your nurse will give you supplies so you can care for your wound at home. Most people also have a nurse visit them at home to help you as you heal. Your case manager (a member of your care team who helps to plan and coordinate your services) will discuss this with you when planning your discharge.

Leaving the hospital

Before you leave the hospital, look at your wound with one of your healthcare providers. Knowing what it looks like will help you notice any changes later.

On the day of your discharge, plan to leave the hospital between 8 a.m. and 10 a.m. Before you leave, one of your healthcare providers will write your discharge order and prescriptions. You'll also get written discharge instructions. One of your healthcare providers will review them with you before you leave.

If your ride isn't at the hospital when you're ready to be discharged, you may be able to wait in the Patient Transition Lounge. A member of your care team will give you more information.

At home

Read *What You Can Do to Avoid Falling* to learn what you can do to keep from falling at home and during your appointments at MSK. You can find it at www.msk.org/pe/avoid_falling or ask for a printed copy.

Filling out your Recovery Tracker

We want to know how you're feeling after you leave the hospital. To help us care for you, we'll send questions to your MSK MyChart account. We'll send them every day for 10 days after you're discharged. These questions are known as your Recovery Tracker.

Fill out your Recovery Tracker every day before midnight (12 a.m.). It only takes 2 to 3 minutes to complete. Your answers to these questions will help us understand how you're feeling and what you need.

Based on your answers, we may reach out to you for more information. Sometimes, we may ask you to call your surgeon's office. You can always contact your surgeon's office if you have any questions.

To learn more, read *Common Questions About MSK's Recovery Tracker*. You can find it at www.msk.org/pe/recovery_tracker or ask for a printed copy.

Caring for your wound

Follow your care team's instructions for how often to change the gauze and bandage over your wound. Most people have a visiting nurse come to their home to help them.

Clean your hands with soap and water or an alcohol-based hand sanitizer before you touch your wound. This will help prevent infection.

Check your wound for signs of infection every day until your healthcare provider tells you it's healed. Call your healthcare provider if you have any of these signs of a wound infection:

- A fever of 100.5 °F (38.0 °C) or higher.
- Swelling around your wound.
- Increased pain that doesn't get better after you take pain medicine.
- Drainage that looks like pus (thick and milky).

Managing your pain

People have pain or discomfort for different lengths of time. You may still have some pain when you go home and will probably be taking pain medicine. The pain usually gets better within 1 to 2 weeks.

Follow these guidelines to help manage your pain at home.

- Take your medicine(s) as directed and as needed.
- Call your healthcare provider if the medicine prescribed for you does not help your pain.
- Do not drive or drink alcohol while you're taking prescription pain medicine. Some prescription pain medicines can make you drowsy (very sleepy). Alcohol can make the drowsiness worse.
- You'll have less pain and need less pain medicine as your incision heals. An over-the-counter pain reliever will help with aches and discomfort. Acetaminophen (Tylenol®) and ibuprofen (Advil® or Motrin®) are examples of over-the-counter pain relievers.
 - Follow your healthcare provider's instructions for stopping your prescription pain medicine.
 - Do not take too much of any medicine. Follow the instructions on the label or from your healthcare provider.
 - Read the labels on all the medicines you're taking. This is very important if you're taking acetaminophen. Acetaminophen is an ingredient in many over-the-counter and prescription medicines. Taking too much can harm your liver. Do not take more than one medicine that has acetaminophen without talking with a member of your care team.
- Pain medicine should help you get back to your usual activities. Take enough to do your activities and exercises comfortably. You may have a little more pain as you start to be more active.
- Keep track of when you take your pain medicine. It works best 30 to 45 minutes after you take it. Taking it when you first have pain is better than waiting for the pain to get worse.

Managing changes in bowel function

Your rectum is a storage tank for stool (poop). You had surgery to remove part of it, so now this tank is smaller.

You may have a lot of small bowel movements because your rectum can't hold as much stool. Over time, your rectum will stretch and be able to hold more. This process can take several months to years.

For the first few weeks after your surgery, you may have a lot of bowel movements. This is because the lower part of your colon hasn't been used in a while. It will take time for your body to recover.

After your surgery, your bowel movements may:

- Be more frequent.
- Happen several times an hour, several times a week.
- Happen every other day.
- Not feel complete. After having a bowel movement, you may still feel like you have to go.

You also may:

- Feel a strong sense of urgency to have a bowel movement.
- Have trouble telling the difference between having a bowel movement and passing gas.

If you also had radiation therapy, your rectum may be stiff. It will not be able to stretch and hold stool as well as before your surgery. This is usually temporary.

Tips for managing frequent bowel movements

If you're having multiple bowel movements a day, you may need to follow a bland diet for a few days. While you're following a bland diet, avoid:

- Dairy products, including milk, cheese and ice cream.
- Coffee.
- Chocolate.
- Spicy foods.
- Fried foods.
- Gravies and cream sauces.
- High-fat deli meats.
- Greasy meats, such as sausage and bacon.
- Fruit juices.
- Sugar-free foods.

Following the BRAT diet can also help control frequent bowel movements. The BRAT diet is made up mostly of:

- Bananas (B)
- White rice (R)
- Applesauce (A)
- Toast (T)

Drinking black tea can also help.

Tips for managing soreness

You may have soreness around your anus from frequent bowel movements.

If you do:

- Soak in warm water 2 to 3 times a day.
- Apply zinc oxide ointment (such as Desitin®) to the skin around your anus after every bowel movement. This helps prevent irritation.
- Don't use harsh toilet paper. You can use a nonalcohol wipe (such as a moistened flushable wipe) instead.
- If your healthcare provider prescribes medicine, take it as directed.

Showering

Taking a warm shower is relaxing and can help decrease muscle aches.

Take your bandages off before you shower. When you shower, use soap to gently wash your wound. After your shower, pat your wound dry with a clean towel. Pack your wound with gauze and change your bandage.

Don't take tub baths or go swimming until your healthcare provider says it's OK.

Eating and drinking

Eat 5 to 6 small meals throughout the day. Remember to eat slowly and chew your food well. It's important you try to maintain your weight. Drink 8 to 10 (8-ounce) cups of liquids every day.

After your surgery, foods may affect you differently. Certain foods may make you have bowel movements right after you eat them. Certain foods may cause diarrhea (loose or watery bowel movements). You may need to change your diet after your surgery.

Every person is different, so there's no way to know which foods will make this happen. During the first few months after your surgery, you'll need to test foods and see how you react to them. It may be helpful to keep a food diary. This will help you keep track of which foods cause discomfort.

If you have questions about your diet, ask to see a clinical dietitian nutritionist.

Dietary supplements and medicines

It may be helpful to take a dietary supplement or medicine to help your bowel function after surgery. **Do not take these things without talking with your healthcare provider first.**

Fiber supplement

Your healthcare provider may tell you to take a fiber supplement (such as Metamucil® or Citrucel®) starting about 4 weeks after your surgery. Fiber will help you by:

- Increasing the bulk in your stool. This helps move waste out of your body because your rectum squeezes better when it's full.
- Stretching your rectum by adding bulk to your stool. This stretching takes place over many months, but eventually your bowel movements should become less frequent and less urgent. This will combine all the small pieces of stool together to create a larger bowel movement. These larger stools will also stretch out your rectum and you will be able to hold more stool.

Start by taking ½ to 1 teaspoon a day and take more of it as needed. Fiber can cause constipation if you're not drinking enough, so drink at least 8 (8-ounce) cups (2 liters) of water or clear liquids each day. If you have bloating and gas, take simethicone (GasX®) as needed.

Probiotic supplement

You may also need to take a probiotic supplement. Taking probiotics may help your bowels work better. Ask your healthcare provider for recommendations.

Other medicines

If you're having many bowel movements, your healthcare provider may prescribe medicine to slow your bowel down. This may help lessen the feeling that you need to go right away and may cut down on your number of bowel movements. Your healthcare provider will tell you how much to take and how often to take it.

If you don't have a bowel movement for 2 days, contact your healthcare provider.

Physical activity and exercise

For the first 6 to 8 weeks after your surgery:

- Don't lift, push, or pull anything heavier than 10 pounds (about 4.5 kilograms).
- Don't do any strenuous activities, such as jogging and tennis.
- Don't play any contact sports, such as football.

Doing aerobic exercise, such as walking and stair climbing, will help you gain strength and feel better. Walk at least 2 to 3 times a day for 15 to 20 minutes for the first 2 weeks after surgery. You can walk outside or indoors at your local mall or shopping center. Ask your healthcare provider before starting more demanding exercises.

It's normal to have less energy than usual after your surgery. Recovery time is different for each person. Increase your activities each day as much as you can. Always balance activity periods with rest periods. But, if you can't sleep at night, it may be a sign that you're resting too much during the day.

Kegel exercises

You'll need to do Kegel exercises to strengthen your pelvic floor muscles. This will help you hold in your bowel movements until you get to a toilet and prevent accidental stool leakage (incontinence).

Ask your healthcare provider when to start doing Kegel exercises. Starting too soon after surgery can be painful.

You can do Kegel exercises while sitting, standing, or walking. To do them:

1. Tighten your pelvic floor muscles. These are the muscles you tighten when you're trying to hold back a bowel movement.
2. Hold this position for 5 to 10 seconds.
3. Release and rest.

Repeat this exercise 10 times, 4 times a day.

Driving

You can start driving again 2 weeks after your surgery. Don't drive while you're taking pain medicine that may make you drowsy. You can ride in a car as a passenger at any time after you leave the hospital.

Going back to work

Talk with your healthcare provider about your job. They'll tell you when it may be safe for you to start working again based on what you do. If you move around a lot or lift heavy objects, you may need to stay out a little longer. If you sit at a desk, you may be able to go back sooner.

Follow-up appointments

Your first appointment after your surgery will be 1 to 3 weeks after you're discharged from the hospital. Call your surgeon's office to schedule it.

It's important to go to all your follow-up appointments after your surgery. You can call your healthcare provider if you have questions between these appointments.

Managing your feelings

You may have new and upsetting feelings after a surgery for a serious illness. Many people say they felt weepy, sad, worried, nervous, irritable, or angry at one time or another. You may find that you cannot control some of these feelings. If this happens, it's a good idea to seek emotional support. Your healthcare provider can refer you to MSK's Counseling Center. You can also reach them by calling 646-888-0200.

The first step in coping is to talk about how you feel. Family and friends can help. We can also reassure, support, and guide you. It's always a good idea to let us know how you, your family, and your friends are feeling emotionally. Many resources are available to you and your family. We're

here to help you and your family and friends handle the emotional aspects of your illness. We can help no matter if you're in the hospital or at home.

When to call your healthcare provider



Call your healthcare provider if:

- You have a fever of 100.5 °F (38.0 °C) or higher.
- You have pain, bloating, cramping, or tenderness in your abdomen.
- You have nausea or vomiting.
- You have trouble passing gas.
- You have trouble having a bowel movement.
- You have trouble urinating (peeing).
- There's swelling around your wound.
- You feel pain around your wound that doesn't go away after taking pain medicine.
- You have bleeding from your rectum.
- You have any of these signs of dehydration (not having enough fluid in your body):
 - Feeling very thirsty.
 - Dry mouth or skin.
 - Fatigue (feeling more tired or weak than usual).
 - Loss of appetite.
 - Feeling dizzy when you stand.
 - Headache.
 - Leg cramps.
- You have any of these signs of wound infection:
 - Swelling.
 - More pain.
 - Warmth at the wound site.
 - Drainage that looks like pus (thick and milky).

Contact information

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call.

If you're not sure how to reach your healthcare provider, call 212-639-2000.

Notes _____

Support services

This section has a list of support services. They may help you as you get ready for your surgery and recover after your surgery.



As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

MSK support services

Admitting Office

212-639-7606

Call if you have questions about your hospital admission, such as asking for a private room.

Anesthesia

212-639-6840

Call if you have questions about anesthesia.

Blood Donor Room

212-639-7643

Call for information if you're interested in donating blood or platelets.

Bobst International Center

www.msk.org/international

888-675-7722

We welcome patients from around the world and offer many services to help. If you're an international patient, call for help arranging your care.

Counseling Center

www.msk.org/counseling

646-888-0200

Many people find that counseling helps them. Our Counseling Center offers counseling for individuals, couples, families, and groups. We can also prescribe medicine to help if you feel anxious or depressed. Ask a member of your care team for a referral or call the number above to make an appointment.

Food Pantry Program

646-888-8055

We give food to people in need during their cancer treatment. Talk with a member of your care team or call the number above to learn more.

Integrative Medicine Service

www.msk.org/integrativemedicine

Our Integrative Medicine Service offers many services to complement (go along with) traditional medical care. For example, we offer music therapy, mind/body therapies, dance and movement therapy, yoga, and touch therapy. Call 646-449-1010 to make an appointment for these services.

You can also schedule a consultation with a healthcare provider in the Integrative Medicine Service. They'll work with you to make a plan for creating a healthy lifestyle and managing side effects. Call 646-608-8550 to make an appointment for a consultation.

MSK Library

library.mskcc.org

212-639-7439

You can visit our library website or call to talk with the library reference staff. They can help you find more information about a type of cancer. You can also visit the library's Patient and Health Care Consumer Education Guide at libguides.mskcc.org/patienteducation

Nutrition Services

www.msk.org/nutrition

212-639-7312

Our Nutrition Service offers nutritional counseling with one of our clinical dietitian nutritionists. Your clinical dietitian nutritionist will talk with you about your eating habits. They can also give advice on what to eat during and after treatment. Ask a member of your care team for a referral or call the number above to make an appointment.

Patient and Community Education

www.msk.org/pe

Visit our patient and community education website to search for educational resources, videos, and online programs.

Patient Billing

646-227-3378

Call if you have questions about preauthorization with your insurance company. This is also called preapproval.

Patient Representative Office

212-639-7202

Call if you have questions about the Health Care Proxy form or concerns about your care.

Perioperative Nurse Liaison

212-639-5935

Call if you have questions about MSK releasing any information while you're having surgery.

Private Duty Nurses and Companions

917-862-6373

You can request private nurses or companions to care for you in the hospital and at home. Call to learn more.

Rehabilitation Services

www.msk.org/rehabilitation

Cancers and cancer treatments can make your body feel weak, stiff, or tight. Some can cause lymphedema (swelling). Our physiatrists (rehabilitation medicine doctors), occupational therapists (OTs), and physical therapists (PTs) can help you get back to your usual activities.

- **Rehabilitation medicine doctors** diagnose and treat problems that affect how you move and do activities. They can design and help coordinate your rehabilitation therapy program, either at MSK or somewhere closer to home. Call Rehabilitation Medicine (Physiatry) at 646-888-1929 to learn more.
- An **OT** can help if you're having trouble doing usual daily activities. For example, they can recommend tools to help make daily tasks easier.

A **PT** can teach you exercises to help build strength and flexibility. Call Rehabilitation Therapy at 646-888-1900 to learn more.

Resources for Life After Cancer (RLAC) Program

646-888-8106

At MSK, care does not end after your treatment. The RLAC Program is for patients and their families who have finished treatment.

This program has many services. We offer seminars, workshops, support groups, and counseling on life after treatment. We can also help with insurance and employment issues.

Sexual Health Programs

Cancer and cancer treatments can affect your sexual health, fertility, or both. MSK's sexual health programs can help you before, during, or after your treatment.

- Our **Female Sexual Medicine and Women's Health Program** can help with sexual health problems such as premature menopause or fertility issues. Ask a member of your MSK care team for a referral or call 646-888-5076 to learn more.
- Our **Male Sexual and Reproductive Medicine Program** can help with sexual health problems such as erectile dysfunction (ED). Ask a member of your care team for a referral or call 646-888-6024 to learn more.

Social Work

www.msk.org/socialwork

212-639-7020

Social workers help patients, families, and friends deal with common issues for people who have cancer. They provide individual counseling and support groups throughout your treatment. They can help you communicate with children and other family members.

Our social workers can also help refer you to community agencies and programs. If you're having trouble paying your bills, they also have information about financial resources. Call the number above to learn more.

Spiritual Care

212-639-5982

Our chaplains (spiritual counselors) are available to listen, help support family members, and pray. They can contact community clergy or faith groups, or simply be a comforting companion and a spiritual presence. Anyone can ask for spiritual support. You do not have to have a religious affiliation (connection to a religion).

MSK's interfaith chapel is located near Memorial Hospital's main lobby. It's open 24 hours a day. If you have an emergency, call 212-639-2000. Ask for the chaplain on call.

Tobacco Treatment Program

www.msk.org/tobacco

212-610-0507

If you want to quit smoking, MSK has specialists who can help. Call to learn more.

Virtual Programs

www.msk.org/vp

We offer online education and support for patients and caregivers. These are live sessions where you can talk or just listen. You can learn about your diagnosis, what to expect during treatment, and how to prepare for your cancer care.

Sessions are private, free, and led by experts. Visit our website to learn more about Virtual Programs or to register.

External support services

There are many other services available to help you before, during, and after your cancer treatment. Some offer support groups and information. Others can help with transportation, lodging, and treatment costs.

Visit www.msk.org/pe/external_support_services for a list of these support services. You can also call 212-639-7020 to talk with an MSK social worker.

Notes _____

Educational resources

This section lists the educational resources mentioned in this guide. It also has copies of the resources that are most important for you to read. They will help you get ready for your surgery and recover after your surgery.



As you read these resources, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

These are the educational resources mentioned in this guide. You can find them online or ask a member of your care team for a printed copy.

- ***A Guide for Caregivers*** (www.msk.org/pe/guide_caregivers)
- ***Advance Care Planning for People With Cancer and Their Loved Ones*** (www.msk.org/pe/advance_care_planning)
- ***Call! Don't Fall!*** (www.msk.org/pe/call_dont_fall)
- ***Common Questions About MSK's Recovery Tracker*** (www.msk.org/pe/recovery_tracker)
- ***Frequently Asked Questions About the Goals to Discharge Checklist*** (www.msk.org/pe/goals_discharge_checklist)
- ***Frequently Asked Questions About Walking After Your Surgery*** (www.msk.org/pe/walking_after_surgery)
- ***Herbal Remedies and Cancer Treatment*** (www.msk.org/pe/herbal_remedies)
- ***How to Be a Health Care Agent*** (www.msk.org/pe/health_care_agent)
- ***How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*** (www.msk.org/pe/check-med-supplement)
- ***How To Use Your Incentive Spirometer*** (www.msk.org/pe/incentive_spirometer)
- ***What You Can Do to Avoid Falling*** (www.msk.org/pe/avoid_falling)

PATIENT & CAREGIVER EDUCATION

Herbal Remedies and Cancer Treatment

This information explains herbal remedies and how they can affect your treatment.

About Herbal Remedies

Herbal remedies are any herbs, botanical (plant-based) supplements, or dietary supplements you take for their health benefits. These may come as tablets, capsules, powders, teas, liquid extracts, and fresh or dried plants.

Some herbal remedies can help prevent or manage side effects of cancer or your treatment. The herbal remedies that can help you depend on what symptoms you have and what treatment you're getting.

Even though herbal remedies can feel safe, they may not all be safe. Herbal remedies do not go through the same testing as prescription medications to make sure they work and are safe.

Some herbal remedies may be harmful. This is because they can:

- Affect how your other medications work.
- Raise or lower your blood pressure.
- Thin your blood and increase your risk of bleeding.
- Keep radiation therapy from working as well as it should.
- Change how your body reacts to sedation (medication to make you calmer) or general anesthesia (medication to make you sleepy).

Talk with your healthcare provider about any herbal remedies or other

supplements you are taking. They can provide an open and safe space to talk about these products.

For more information about herbs and supplements, visit www.aboutherbbs.com or call MSK's Integrative Medicine Service at 646-608-8550.

Stop taking herbal remedies before your treatment

Stop taking herbal remedies and other dietary supplements 7 days (1 week) before you:

- Have surgery.
- Start chemotherapy.
- Start radiation therapy.
- Have certain procedures. Your healthcare provider will let you know if you need to stop taking herbal remedies before your procedure.

Herbal remedies and other dietary supplements can cause bleeding and affect your treatment. Follow your healthcare provider's instructions for when to restart taking herbal remedies.

You can still use some herbs in your food and drinks, such as using spices in cooking and drinking tea. Herbal remedies are stronger than the herbs you cook with.

Common Herbal Remedies and Their Effects

These are some commonly used herbs and their side effects on cancer treatments.

Echinacea (EH-kih-NAY-shuh)

- Can cause rare but serious allergic reactions, such as a rash or trouble breathing.
- Can keep medications that weaken your immune system from working as well as they should.

Garlic

- Can lower your blood pressure and cholesterol levels.
- Can increase your risk of bleeding.

Ginkgo (also known as Ginkgo biloba)

- Can increase your risk of bleeding.

Ginseng (JIN-seng)

- Can keep sedation or general anesthesia from working as well as they should.
- Can increase your blood pressure.
- Can increase your risk of bleeding.
- Can lower your blood glucose (sugar) level.

Turmeric (TER-mayr-ik)

- Can keep chemotherapy from working as well as it should.

St. John's Wort

- Can keep some medications from working as well as they should.
- Can make your skin more sensitive to radiation or laser treatment.

Valerian (vuh-LEER-ee-un)

- Can make sedation or general anesthesia affect you more than they should.

Herbal formulas

- Herbal formulas contain many different herbs and dosages.
- Stop taking these products 7 days (1 week) before treatment. Do not start taking herbal formulas again until your healthcare provider tells you it is safe.

This information does not cover all herbal remedies or possible side effects. Talk with your healthcare provider if you have any questions or concerns.

Contact Information

- To schedule a consultation with a healthcare provider in Integrative Medicine, call 646-608-8550.
- To make an appointment for Integrative Medicine Service's therapies, classes, and workshops, call 646-449-1010.

For more information, visit www.mskcc.org/IntegrativeMedicine or read *Integrative Medicine Therapies and Your Cancer Treatment* (www.mskcc.org/pe/integrative_therapies).

For more resources, visit www.mskcc.org/pe to search our virtual library.

Herbal Remedies and Cancer Treatment - Last updated on May 5, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil

This information will help you check if your medicines or dietary supplements have aspirin, other NSAIDs, vitamin E, or fish oil as an active ingredient. NSAID stands for nonsteroidal anti-inflammatory drug.

It's important to stop taking these medicines and supplements before many cancer treatments. They affect your platelets (blood cells that clot to prevent bleeding) and can raise your risk of bleeding.

Other dietary supplements, such as vitamins and herbal remedies, can also affect your cancer treatment. Read *Herbal Remedies and Cancer Treatment* (www.mskcc.org/pe/herbal_remedies) to learn more.

Make sure your healthcare provider always knows all the prescription and over-the-counter medicines and supplements you're taking. This includes patches and creams.

A prescription medicine is one you can only get with a prescription from your healthcare provider. An over-the-counter medicine is one you can buy without a prescription.

What is an active ingredient?

An active ingredient is the part of a medicine or supplement that makes it work. Some medicines and supplements have just one active ingredient. Others have more. For example:

- Ibuprofen is the active ingredient in Advil® and Motrin®. Ibuprofen is an NSAID.
- Naproxen is the active ingredient in Aleve®. Naproxen is an NSAID.
- Acetaminophen is the active ingredient in Tylenol®.
- Aspirin, acetaminophen, and caffeine are the active ingredients in Excedrin®.

Generic medicines sometimes use their active ingredient as their name. But people often call medicines and supplements by a brand name, even if they're generic. This can make it hard to know their active ingredients.

How to find a medicine or supplement's active ingredients

You can always find the active ingredients by reading the label.

Over-the-counter medicines

Over-the-counter medicines list their active ingredients in the “Drug Facts” label (see Figure 1). Active ingredients are always the first thing on the Drug Facts label.



Figure 1. Active ingredients on an over-the-counter medicine label

Prescription medicines

Prescription medicines list their active ingredients on the label. Their active ingredients and their generic name are the same thing.

Labels often look different depending on which pharmacy you use. Here's an example of where to find a medicine's active ingredients (generic name) on a label from MSK's pharmacy (see Figure 2).




Figure 2. Active ingredients on a prescription medicine label

Dietary supplements

Dietary supplements list their active ingredients in the "Supplement Facts" label (see Figure 3). The active ingredients always have an amount per serving and % daily value included.

Supplement Facts		
Serving Size 1 Tablet		
	Amount Per Serving	% Daily Value
Vitamin A (as retinyl acetate and 50% as beta-carotene)	5000 IU	100%
Vitamin C (as ascorbic acid)	60 mg	100%
Vitamin D (as cholecalciferol)	400 IU	100%
Vitamin E (as di-alpha tocopheryl acetate)	30 IU	100%
Thiamin (as thiamin mononitrate)	1.5 mg	100%
Riboflavin	1.7 mg	100%
Niacin (as niacinamide)	20 mg	100%
Vitamin B ₆ (as pyridoxine hydrochloride)	2.0 mg	100%
Folate (as folic acid)	400 mcg	100%
Vitamin B ₁₂ (as cyanocobalamin)	6 mcg	100%
Biotin	30 mcg	10%
Pantothenic Acid (as calcium pantothenate)	10 mg	100%
Other ingredients: Gelatin, lactose, magnesium stearate, microcrystalline cellulose, FD&C Yellow No. 6, propylene glycol, propylparaben, and sodium benzoate.		

Figure 3. Active ingredients on a supplement label

Active ingredients to look for

If your medicine or supplement has any of these active ingredients, you may need to stop taking it before, during, or after your cancer treatment or surgery. Follow your care team's instructions.

Active ingredients to look for		
<ul style="list-style-type: none">• Acetylsalicylic acid• Alpha-linolenic acid (ALA)• Aspirin• Acetaminophen*• Celecoxib• Diclofenac• Diflunisal• Docosahexaenoic acid (DHA)• Eicosapentaenoic acid (EPA)	<ul style="list-style-type: none">• Etodolac• Fish oil• Fenoprofen Flurbiprofen• Ibuprofen• Indomethacin• Ketoprofen• Ketorolac• Meclofenamate• Mefenamic acid• Meloxicam	<ul style="list-style-type: none">• Nabumetone• Naproxen• Omega-3 fatty acids• Omega-6 fatty acids• Oxaprozin• Piroxicam• Sulindac• Tolmetin• Vitamin E

* The full name acetaminophen isn't always written out. Look for the common abbreviations listed below, especially on prescription pain relievers.

Common abbreviations for acetaminophen		
<ul style="list-style-type: none">• APAP• Acetamin	<ul style="list-style-type: none">• AC• Acetam	<ul style="list-style-type: none">• Acetaminop• Acetaminoph

About acetaminophen (Tylenol)

In general, acetaminophen is safe to take during cancer treatment. It doesn't affect platelets. That means it will not raise your chance of bleeding. If you're getting chemotherapy, talk with your healthcare provider before taking acetaminophen.

There is a limit to how much acetaminophen you can take in a day. Always follow the instructions from your care team or on the medicine's label.

Acetaminophen is in many different prescription and over-the-counter medicines. It's possible to take too much without knowing. **Always read the label on the medicines you take.** Do not take more than 1 medicine that has acetaminophen at a time without talking with a member of your care team.

Instructions before your cancer treatment

Tell your healthcare provider if you take aspirin, other NSAIDs, vitamin E, or fish oil. They'll tell you if you need to stop taking it. You'll also find instructions in the information about your treatment.

Before your surgery

Follow these instructions if you're having surgery or a surgical procedure. **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take vitamin E, fish oil, or a supplement that has vitamin E or fish oil, stop taking it 7 days before your surgery or as directed by your healthcare provider.
- If you take an NSAID or a medicine that has an NSAID, stop taking it 48 hours (2 days) before your surgery or as directed by your healthcare provider.

Before your radiology procedure

Follow these instructions if you're having a radiology procedure (including Interventional Radiology, Interventional Mammography, Breast Imaging, and General Radiology). **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to stop taking it 5 days before your procedure. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take an NSAID or a medicine that has an NSAID, you may need to stop taking it 24 hours (1 day) before your procedure. Follow your healthcare provider's instructions.

Before and during your chemotherapy

Chemotherapy can lower your platelet count, which can increase your risk of bleeding. No matter if you're just starting chemotherapy or have been getting it, talk with your healthcare provider before taking aspirin, other NSAIDs, vitamin E, or fish oil.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil - Last updated on November 29, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center