

# How To Do Bowel Prep Before Your Surgery at MSK Kids

This information explains how to do bowel preparation (bowel prep) before your surgery at MSK Kids. Doing bowel prep helps clear stool (poop) from your body.

In this resource, the words “you” and “your” mean you or your child.

## Bowel prep supplies

You can buy these at your local pharmacy without a prescription.

- Polyethylene glycol (MiraLAX®) powder
- Clear liquids that are not red, orange, or purple

Read the section “Get the MiraLAX mixture ready” to see how much you’ll need. Make sure to get enough clear liquids to have while you’re following a clear liquid diet.

Your healthcare provider will also give you a prescription for the medicines below. Make sure you have them before you start your bowel prep.

- Metronidazole
- Neomycin

# Bowel prep instructions

Start your bowel prep the day before your surgery.

## Get the MiraLAX mixture ready

Mix \_\_\_\_\_ grams of MiraLAX with \_\_\_\_\_ ounces of a clear liquid in the morning the day before your surgery. Stir the mixture until the MiraLAX powder dissolves. Once the powder dissolves, you can put the mixture in the refrigerator.

## Start following a clear liquid diet at noon (12 p.m.)

Start following a clear liquid diet at noon (12 p.m.) the day before your surgery.

A clear liquid diet includes only liquids you can see through. You can find examples in the “Clear liquid diet” table. Avoid clear liquids that are red, orange, or purple.

While you’re following a clear liquid diet:

- Do not eat any solid foods.
- Drink different types of clear liquids. Do not just drink water, coffee, and tea. This helps make sure you get enough calories and is an important part of your bowel prep.
- Do not drink any liquids you cannot see through, such as milk or smoothies.
- Do not drink sugar-free liquids unless you have diabetes and your healthcare provider tells you to.

## Clear liquid diet

	OK to have	Do not have
Soups	<ul style="list-style-type: none"><li>Clear broth, bouillon, or consommé.</li></ul>	<ul style="list-style-type: none"><li>Anything with pieces of food or seasoning.</li></ul>
Sweets	<ul style="list-style-type: none"><li>Gelatin, such as Jell-O®.</li><li>Flavored ice.</li><li>Hard candies, such as Life Savers®, lemon drops, and peppermints.</li></ul>	<ul style="list-style-type: none"><li>Anything red, orange, or purple.</li><li>All other sweets.</li></ul>
Drinks	<ul style="list-style-type: none"><li>Clear fruit juices, such as lemonade, apple, white cranberry, and white grape juices.</li><li>Soda, such as ginger ale, 7UP®, Sprite®, and seltzer.</li><li>Sports drinks, such as Gatorade® and Powerade®.</li><li>Coffee without milk or creamer.</li><li>Tea without milk or creamer.</li><li>Water, including carbonated (fizzy) and flavored water.</li><li>Clear nutritional drinks, such as Boost® Breeze, Ensure Clear™, Pedialyte®, and Diabetishield®.</li></ul>	<ul style="list-style-type: none"><li>Anything red, orange, or purple</li><li>Juices with pulp.</li><li>Nectars.</li><li>Smoothies or shakes.</li><li>Milk, cream, and other dairy products.</li><li>Nut milks, plant milks, non-dairy creamers, and other dairy alternatives.</li><li>Drinks with alcohol.</li></ul>

## Start drinking the MiraLAX mixture at 2 p.m.

Start drinking the MiraLAX mixture at 2 p.m. the day before your surgery.

Drink the MiraLAX mixture over the next 3 to 4 hours. Aim to finish between 5 p.m. and 6 p.m.

The MiraLAX will make you have bowel movements often. Make sure you're near a bathroom once you start taking it.

Having lots of bowel movements can irritate (bother) your anus (where your poop leaves your body). Putting petroleum jelly (Vaseline®) or A & D® ointment on the skin around your anus after each bowel movement can help.

## Take neomycin and metronidazole at 6 p.m. and 10 p.m.

Take \_\_\_\_\_ (1 dose) of neomycin and \_\_\_\_\_ (1 dose) of metronidazole at 6 p.m. the night before your surgery.

Take another \_\_\_\_\_ (1 dose) of neomycin and \_\_\_\_\_ (1 dose) of metronidazole at 10 p.m. the night before your surgery.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

How To Do Bowel Prep Before Your Surgery at MSK Kids - Last updated on February 21, 2024

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center