



---

## PATIENT & CAREGIVER EDUCATION

# Entrectinib

This information from Lexicomp<sup>®</sup> explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### **Brand Names: US**

Rozlytrek

### **Brand Names: Canada**

Rozlytrek

### **What is this drug used for?**

- It is used to treat cancer.

### **What do I need to tell my doctor BEFORE I take this drug?**

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you take any other drugs (prescription or OTC, natural products, vitamins). There are many drugs that interact with this drug, like certain drugs that are

used for HIV, infections, or seizures.

- If you are taking any drugs that can cause a certain type of heartbeat that is not normal (prolonged QT interval). There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 1 week after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

### **What are some things I need to know or do while I take this drug?**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.

- Have your blood work and heart function checked as you have been told by your doctor.
- Avoid grapefruit and grapefruit juice.
- If you have gout, talk with your doctor.
- High uric acid levels have happened with this drug. Call your doctor right away if your big toe or any other joint is painful, red, hot, tender, stiff, or swollen.
- Heart failure has happened with this drug, as well as heart failure that has gotten worse in people who already have it. Tell your doctor if you have heart disease. Call your doctor right away if you have shortness of breath, a big weight gain, a heartbeat that is not normal, or swelling in the arms or legs that is new or worse.
- This drug may raise the chance of a broken bone. Talk with the doctor.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.
- This drug may cause harm to an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.
- If you or your sex partner may become pregnant, you must use birth control while taking this drug and for some time after the last dose. Ask your doctor how

long to use birth control. If you or your sex partner gets pregnant, call your doctor right away.

## **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of nervous system problems like a burning, numbness, or tingling feeling; change in balance; confusion; dizziness; hallucinations; memory problems or loss; new or worse behavior or mood changes like anxiety, depression, or thoughts of suicide; trouble focusing; trouble sleeping; or trouble thinking or speaking.
- Signs of liver problems like dark urine, tiredness,

decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Bone pain.
- Change in eyesight.
- If bright lights bother your eyes.
- Swelling.
- A type of abnormal heartbeat (prolonged QT interval) can happen with this drug. Call your doctor right away if you have a fast heartbeat, a heartbeat that does not feel normal, or if you pass out.
- Low white blood cell counts have happened with this drug. This may lead to a higher chance of getting an

infection. Call your doctor right away if you have signs of infection like fever, chills, or sore throat.

### **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Change in taste.
- Muscle pain or weakness.
- Back pain.
- Pain in arms or legs.
- Weight gain.
- Cough.
- Headache.
- Stuffy nose.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

### **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

### **All products:**

- Take with or without food.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- If you throw up right after taking this drug, take 1 more dose as soon as you can on the same day. Then go back to your normal time the next day.

### **Capsules:**

- Swallow whole. Do not chew, break, or dissolve.
- A liquid (suspension) can be made from the capsules if needed. Talk with the doctor or pharmacist.
- If you make a suspension, take this drug right after mixing it. Drink water after taking this drug to make

sure the dose is fully swallowed. Throw away any part of the suspension not used within 2 hours after mixing.

- Those who have feeding tubes may use this drug. Use as you have been told. Flush the feeding tube after this drug is given.

### **Oral pellets:**

- Sprinkle the pellets on 1 or more spoonfuls of a soft food like applesauce, yogurt, or pudding. Do not crush or chew. Take within 20 minutes after putting this drug on the food. Be sure you take the whole dose of this drug.
- Throw away the soft food with the pellets sprinkled on it if not taken within 20 minutes and make a new dose.
- Drink water after you take this drug to make sure the dose is fully swallowed.
- Use the whole packet of pellets. Do not try to measure out only some of the pellets from a packet for a dose.
- Do not put this form of this drug down a feeding tube.
- Do not use pellets to make a suspension.



## **What do I do if I miss a dose?**

- Take a missed dose as soon as you think about it.
- If it is less than 12 hours until the next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

## **How do I store and/or throw out this drug?**

- Store in the original container at room temperature.
- Keep lid tightly closed.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## **General drug facts**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information

leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

### **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients

must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2024-02-05

### **Copyright**

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Entrectinib - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center