

#### PATIENT & CAREGIVER EDUCATION

# Famotidine

This information from Lexicomp<sup>®</sup> explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

# **Brand Names: US**

Acid Reducer Maximum Strength [OTC]; Acid Reducer [OTC]; Famotidine Maximum Strength [OTC]; Famotidine Orig St [OTC]; FT Acid Reducer Max Strength [OTC]; FT Acid Reducer [OTC]; Heartburn Relief Max St [OTC]; Heartburn Relief [OTC]; Pepcid; Pepcid AC Maximum Strength [OTC]; Pepcid AC [OTC]

# **Brand Names: Canada**

AG-Famotidine; ALTI-Famotidine; CO Famotidine; Famotidine Omega W-O Preserv; Famotidine Omega W-Preserv; GMD-Famotidine; JAMP Famotidine; MINT-Famotidine; TEVA-Famotidine

# What is this drug used for?

- It is used to treat or prevent GI (gastrointestinal) ulcers.
- It is used to treat gastroesophageal reflux disease (GERD; acid reflux).
- It is used to treat heartburn and sour stomach.
- It is used to treat syndromes caused by lots of stomach acid.
- It may be given to you for other reasons. Talk with the doctor.

# What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Black or bloody stools; heartburn with light-headedness, sweating, dizziness, or wheezing; chest pain; shoulder pain with shortness of breath; pain that spreads to the arms, neck, or shoulders; lightheadedness; sweating a lot; throwing up blood; or trouble or pain swallowing food.
- If you have had heartburn for 3 months or more.
- If you are taking any of these drugs: Cefditoren, dasatinib, delavirdine, or fosamprenavir.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

# What are some things I need to know or do while I take this drug? All products:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Do not take this drug for longer than you were told by your doctor.
- If you have kidney problems or are 65 or older, talk with your doctor. Hallucinations (seeing or hearing things that are not there); seizures; and feeling agitated, confused, sluggish, or out of sorts have happened in people with kidney problems and in older people.

- This drug may prevent other drugs taken by mouth from getting into the body. If you take other drugs by mouth, you may need to take them at some other time than this drug. Talk with your doctor.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

#### Injection:

• Some products have benzyl alcohol. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol. Talk with the doctor to see if this product has benzyl alcohol in it.

# What are some side effects that I need to call my doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Dizziness or passing out.
- Fast or abnormal heartbeat.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

• Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

### How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

#### All oral products:

- Take with or without food.
- Ask your doctor before you take antacids with this drug.

#### **Tablets:**

- Do not chew this drug.
- To prevent heartburn, take before eating foods or drinking liquids that cause heartburn. Take as you were told by your doctor or on the package labeling.

#### **Chewable tablets:**

• Chew or crush well. Do not swallow it whole.

## Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

#### Injection:

• It is given as a shot into a vein.

### What do I do if I miss a dose? All oral products:

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- Many times this drug is taken on an as needed basis. Do not take more often than told by the doctor.

#### Injection:

• Call your doctor to find out what to do.

#### How do I store and/or throw out this drug? All oral products:

• Store at room temperature in a dry place. Do not store in a bathroom.

#### Liquid (suspension):

• Throw away any part not used after 1 month.

#### Injection:

• If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

#### All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour

down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## **General drug facts**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

# **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

### **Last Reviewed Date**

2023-01-25

# Copyright

@ 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Famotidine - Last updated on December 12, 2022 All rights owned and reserved by Memorial Sloan Kettering Cancer Center