



PATIENT & CAREGIVER EDUCATION

Leuprolide

This information from Lexicomp[®] explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Camcevi; Eligard; Fensolvi (6 Month); Lupron Depot (1-Month); Lupron Depot (3-Month); Lupron Depot (4-Month); Lupron Depot (6-Month); Lupron Depot-Ped (1-Month); Lupron Depot-Ped (3-Month); Lupron Depot-Ped (6-Month)

Brand Names: Canada

Eligard; Lupron Depot; Lupron Depot (1-Month); Lupron Depot (3-Month); Lupron Depot 3 Month Kit; Lupron [DSC]; Zeulide Depot

What is this drug used for?

- It is used to treat endometriosis.
- It is used to treat anemia caused by fibroids of the uterus.

- It is used to treat prostate cancer.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

For all patients taking this drug:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

Females:

- If you have unexplained vaginal bleeding.
- If you are pregnant or may be pregnant. Do not take this drug if you are pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

For all patients taking this drug:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- This drug may raise some hormone levels in your body during the first few weeks of taking it. Disease signs may get worse before getting better. Tell your doctor if you have any new signs or if your disease signs are worse for longer than a few weeks after starting this drug.
- This drug lowers some hormone levels in your body. This may cause some effects like change in breast size, breast soreness or tenderness, testicle changes, trouble getting or keeping an erection, lowered interest in sex, hot flashes, or sweating. Talk with your doctor.
- Have your blood work and bone density checked as you have been told by your doctor.
- High blood sugar has happened with this drug. This includes diabetes that is new or worse.
- Check your blood sugar as you have been told by your doctor.

- If you smoke, talk with your doctor.
- High cholesterol has happened with this drug. If you have questions, talk with the doctor.
- A very bad pituitary gland problem (pituitary apoplexy) has rarely happened with this drug. Most of the time, this has happened within 2 weeks after the first dose. Call your doctor right away if you have a sudden headache, throwing up, passing out, mood changes, eye weakness, not able to move your eyes, or change in eyesight.
- This drug may cause weak bones. This may happen more often if used for a long time. This may raise the chance of broken bones. Call your doctor right away if you have bone pain.
- A higher chance of heart attack, stroke, or sudden heart death has been noted when drugs like this were used in males. The chance is low, but get medical help right away if you have chest pain or pressure, weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on 1 side of the face, or change in eyesight.
- Some products are not approved for use in people who are 65 or older or in children. Talk with your doctor.

Males:

- The chance of severe and sometimes deadly problems may be raised in people with bladder blockage. It may also be raised in people with growths on or near the spine or spinal cord. If you have questions, talk with your doctor.
- Lowering male hormones in the body may raise the chance of a type of heartbeat that is not normal called prolonged QT interval. Talk with the doctor.
- This drug may affect being able to father a child. Talk with the doctor.

Females:

- This drug may affect being able to get pregnant. This effect goes back to normal when the drug is stopped. If you have questions, talk with the doctor.
- Most of the time, this drug stops you from having a period (menstrual bleeding). This is not a method of birth control. Use a non-hormone type of birth control like condoms to prevent pregnancy while taking this drug.
- If you miss doses of this drug, bleeding between cycles can happen. Talk with your doctor.
- This drug may cause harm to the unborn baby if you

take it while you are pregnant.

- A pregnancy test will be done to show that you are NOT pregnant before starting this drug. If you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

For all patients taking this drug:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more

often, flushing, fast breathing, or breath that smells like fruit.

- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of dehydration like dry skin, mouth, or eyes; thirst; fast heartbeat; dizziness; fast breathing; or confusion.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Bone pain that is new or worse.
- Not able to pass urine or change in how much urine is passed.
- Very bad back pain.
- Seizures.
- Blood in the urine.
- A burning, numbness, or tingling feeling that is not normal.

- Severe dizziness or passing out.
- Fast, slow, or abnormal heartbeat.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Not able to move.
- Behavior and mood changes have happened with the use of drugs like this one in children. This includes acting aggressive, crying, depression, emotional ups and downs, restlessness, and feeling angry and irritable. Call your doctor right away if you have any new or worse behavior or mood changes.
- A severe skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause severe health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes.

Females:

- Still having a menstrual period.
- Vaginal itching or discharge.
- Vaginal irritation.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

For all patients taking this drug:

- Headache.
- Feeling dizzy, tired, or weak.
- Decreased appetite.
- Constipation.
- Upset stomach or throwing up.
- Trouble sleeping.
- Irritation where the shot is given.
- Back, muscle, or joint pain.
- Flu-like signs.
- Signs of a common cold.

Females:

- Pimples (acne).
- Mood changes.
- A change in weight without trying.

- Feeling nervous and excitable.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Some products need to be given into the fatty part of the skin. Some products need to be given into a muscle. Talk with the doctor or pharmacist if you are not sure how to use this drug.
- Wash your hands before and after use.
- Wear gloves while mixing and giving this drug.
- Move the site where you give the shot with each shot.

- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.
- Keep using this drug as you have been told by your doctor or other health care provider, even if you feel well.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- Most of the time, this drug will be given in a hospital or doctor's office. If stored at home, follow how to store as you were told by the doctor.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush

down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-12-27

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Leuprolide - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center