



PATIENT & CAREGIVER EDUCATION

Methyl Aminolevulinate

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: Canada

Metvix

Warning

- Only put this drug on the areas you were told. Keep this drug out of your eyes, nose, mouth, vagina, and rectum.
- Do not put on healthy skin.
- If you get this drug in any of these areas, rinse well with water.

What is this drug used for?

- It is used to treat a precancerous skin problem called actinic keratosis.
- It is used to treat a type of skin cancer called basal cell carcinoma.

What do I need to tell my doctor **BEFORE** I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

- If you are allergic to almond oil, peanuts, or porphyrins.
- If your skin is sensitive to light.
- If you have porphyria.
- If you have a type of skin cancer called morpheaform basal cell carcinoma.
- If you are taking any drugs that may make your skin more sensitive to light. There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.
- If you are breast-feeding. Do not breast-feed for at least 48 hours after getting this drug.
- If the patient is a child. Do not give this drug to a child.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- The treated area may be more sensitive to light. Light may cause a burning or stinging feeling. Protect the treated area and skin around it for at least 48 hours after you get this drug and have light therapy. Protect it from sun, sunlamps, bright indoor lights, and tanning beds. Wear the proper covering like a wide-brimmed hat, long sleeve shirt, or gloves to protect the treated skin from light. Sunscreens will not help.
- The treated area may be more sensitive to light. Light may cause a burning or stinging feeling. Protect the treated area and skin around it

from sun, sunlamps, bright indoor lights, and tanning beds after this drug is put on and for as long as you have been told by your doctor. Wear the proper covering like a wide-brimmed hat, long sleeve shirt, or gloves to protect the treated skin from light. Sunscreens will not help.

- Do not use sunscreen or other drugs on affected part.
- Protect from cold.
- Talk with your doctor before you use other drugs or products on your skin. This includes soaps.
- Call your doctor right away if you have a skin reaction that bothers you or if the reaction makes it hard to keep putting this drug on. Call your doctor right away if you have a skin reaction that causes problems with daily living.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Any unexplained bruising or bleeding.
- Redness, swelling, or other skin irritation that gets worse or lasts longer than 3 weeks.
- High blood pressure caused by pain from the procedure has happened. If you have pain from the procedure, you may need to have your blood

pressure checked. Tell your doctor right away if you have signs of high blood pressure like severe headache or dizziness, passing out, or change in eyesight.

- Rarely, short-term memory loss or confusion have happened after light therapy. Tell your doctor right away if you have these effects.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Headache.
- Upset stomach.
- Feeling tired.
- People using this drug may have skin reactions where this drug is used. These reactions may include bleeding, blisters, burning, crusting, itching, pain, peeling, redness, stinging, swelling, or ulcers. Most of the time, burning, pain, and stinging go away within a few hours. If these reactions get very bad, a break from using this drug may be needed as told by the doctor. Talk with the doctor.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Your doctor or other healthcare provider will put on the skin.
- This drug is used with light therapy.
- You will need to protect your eyes during the light therapy. Follow what your doctor has told you.
- If you cannot have the light therapy after this drug has been put on, talk with your doctor. Avoid bright light for as long as you have been told by your doctor.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- This drug will be given in a healthcare setting. You will not store it at home.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk

with your doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-03-22

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Methyl Aminolevulinate - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center