



---

PATIENT & CAREGIVER EDUCATION

# Omeprazole, Amoxicillin, and Rifabutin

This information from Lexicomp<sup>®</sup> explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## **Brand Names: US**

Talicia

## **What is this drug used for?**

- It is used to treat GI (gastrointestinal) ulcers caused by infection.

## **What do I need to tell my doctor BEFORE I take this drug?**

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have mono.
- If you have any of these health problems: Kidney disease or liver disease.

- If you have had kidney problems caused by this drug or a drug like this one.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, seizures, and others. There are many drugs that must not be taken with this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

### **What are some things I need to know or do while I take this drug?**

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.

- This drug may stain contact lenses.
- This drug may affect being able to father a child. Talk with the doctor.
- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Use some other kind of birth control also like a condom when taking this drug.
- This drug may cause harm to the unborn baby if you take it while you are pregnant. If you are pregnant or you get pregnant while taking this drug, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

### **What are some side effects that I need to call my doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat;

trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been deadly.

- Signs of a type of allergic reaction called drug-induced enterocolitis syndrome, like vomiting within 1 to 4 hours after taking this drug, diarrhea within 24 hours after taking this drug, pale or gray skin, feeling tired or unwell, or signs of low blood pressure like severe dizziness or passing out.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Change in eyesight, eye pain, or severe eye irritation.
- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.
- Flu-like signs. These include headache, weakness, fever, shakes, aches, pains, and sweating.
- Chest pain or pressure.
- A heartbeat that does not feel normal.
- Shortness of breath.
- Muscle or joint pain.
- Swelling.

- Cough.
- Diarrhea is common with antibiotics. Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem. CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea.
- Severe skin reactions may happen with this drug. These include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be affected. These reactions can be deadly. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in your mouth, throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.
- Lupus has happened with this drug, as well as lupus that has gotten worse in people who already have it. Tell your doctor if you have lupus. Call your doctor right away if you have signs of lupus like a rash on the cheeks or other body parts, change in skin color, sunburn easy, muscle or joint pain, chest pain or

shortness of breath, or swelling in the arms or legs.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Diarrhea.
- Headache.
- This drug may change the color of urine, stools, saliva, sweat, tears, and skin to a brown-orange color. This is normal and not harmful.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

## **How is this drug best taken?**

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take this drug with food.
- Swallow whole with a full glass of water.

- Do not chew or crush.
- Do not drink alcohol at the time you take this drug.
- Use as you have been told, even if your signs get better.

### **What do I do if I miss a dose?**

- If it is more than 4 hours until your next dose, take the missed dose as soon as you think about it and go back to your normal time.
- If it is less than 4 hours until your next dose, take the missed dose as soon as you think about it. Wait to take your next dose until at least 4 hours after you take the missed dose.
- Do not take 2 doses at the same time or extra doses.
- If you are not sure what to do if you miss a dose, call your doctor.

### **How do I store and/or throw out this drug?**

- Store in the original container at room temperature.
- Keep lid tightly closed.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush

down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## **General drug facts**

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments,



medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2024-06-05

### **Copyright**

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Omeprazole, Amoxicillin, and Rifabutin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center