



PATIENT & CAREGIVER EDUCATION

Pemigatinib

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Pemazyre

Brand Names: Canada

Pemazyre

What is this drug used for?

- It is used to treat bile duct cancer.
- It is used to treat a type of blood cancer called myeloid/lymphoid neoplasms (MLNs).

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you take any other drugs (prescription or OTC, natural products, vitamins). There are many drugs that interact with this drug, like certain drugs that are used for HIV, infections, or seizures.
- If you are breast-feeding. Do not breast-feed while you take this drug and for 1 week after your last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have your blood work checked and eye exams as you have been told by your doctor.
- It is common to have dry eyes with this drug. Artificial tears, lubricant eye gels, or other products may be used to prevent or treat dry eyes. If you have questions, talk with the doctor or pharmacist.
- If you have upset stomach, throwing up, diarrhea, or decreased appetite, talk with your doctor. There may be ways to lower these side effects.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- Tell your doctor if you have signs of high or low blood sugar like breath that smells like fruit, dizziness, fast breathing, fast heartbeat, feeling confused, feeling sleepy, feeling weak, flushing, headache, unusual thirst or hunger, passing urine more often, shaking, or sweating.
- Avoid grapefruit and grapefruit juice.
- It is common to have high phosphate levels in the blood with this drug. Sometimes, this can be severe and can lead to certain health problems. If you have questions, talk with the doctor.
- This drug may cause harm to an unborn baby or loss of an unborn baby. A pregnancy test will be done before you start this drug to show that you are NOT pregnant.

- If you or your sex partner may become pregnant, you must use birth control while taking this drug and for some time after the last dose. Ask your doctor how long to use birth control. If you or your sex partner gets pregnant, call your doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Redness or irritation of the palms of hands or soles of feet.
- Swelling in the arms or legs.
- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.
- Numbness or tingling around the mouth.

- Rarely, severe eye problems (retinal tear or detachment) have happened with this drug. Call your doctor or get medical help right away if you start to see flashing lights, floaters, a curtain-like shadow coming across your eye, or have sudden eyesight loss.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Hair loss.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Dry mouth, skin, or eyes.
- More tears.
- Eyelash changes.
- Change in nails.
- Mouth irritation or mouth sores.
- Feeling dizzy, tired, or weak.
- Change in taste.
- Headache.
- Back, joint, arm, or leg pain.
- Weight loss.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food.
- Swallow whole. Do not chew, break, crush, or dissolve.
- Take this drug at the same time of day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- If you throw up after taking a dose, do not repeat the dose. Take your next dose at your normal time.

For patients with bile duct cancer:

- There may be days when you will not take this drug.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it has been 4 hours or more since the missed dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2022-09-09

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Pemigatinib - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center