



PATIENT & CAREGIVER EDUCATION

Simeprevir

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Warning

- You will be tested for hepatitis B before starting this drug. In people who have had hepatitis B or carry the virus, the virus has become active again during or after treatment with drugs like this one. This can lead to severe and sometimes deadly liver problems. People with hepatitis B and hepatitis C will be watched closely during and after treatment.

What is this drug used for?

- It is used to treat long-term hepatitis C infections.

What do I need to tell my doctor **BEFORE** I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have ever had liver problems other than hepatitis C.
- If you are **not** taking other drugs to treat hepatitis C.
- If you take any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, seizures, and others. There are many drugs that must not be taken with this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- If you have a sulfa (sulfonamide) allergy, talk with your doctor.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- Avoid grapefruit and grapefruit juice.
- You may get sunburned more easily. Avoid sun, sunlamps, and tanning beds. Use sunscreen and wear clothing and eyewear that protects you from the sun.
- Very bad skin reactions to sunlight have happened with this drug. Sometimes, this has led to need to go to the hospital. Talk with the doctor.
- Liver problems have happened with this drug. Liver problems have also gotten worse in people who already have them. Rarely, this has been deadly. If you have questions, talk with the doctor.
- Slow heartbeat and the need to get a pacemaker have happened when amiodarone was given with sofosbuvir and certain other hepatitis C drugs. Sometimes, this has been deadly. Slow heartbeat has happened up to 2

weeks after starting hepatitis C treatment. You will need to be watched closely if you will be taking amiodarone with hepatitis C treatment. Follow what your doctor has told you to do. Call your doctor right away if you have signs of slow heartbeat like chest pain, confusion, dizziness, passing out or near-passing out, memory problems, shortness of breath, tiredness, or weakness.

- This drug does not stop the spread of hepatitis to others through blood, body fluids, or having sex. Do not have any kind of sex without using a condom. Do not share injection equipment like needles. Do not share personal items that can have blood or body fluids on them like toothbrushes or razors. Talk with your doctor or other health care provider about safe sex and other ways to prevent spreading hepatitis to others.
- This drug must be taken with other drugs to treat hepatitis C. Be sure you know about the warnings, benefits, and risks of the other drugs. Talk with your doctor if you have questions or concerns about any of your drugs used to treat hepatitis C.
- If you stop taking any of your other drugs to treat hepatitis C, talk with your doctor. You may need to stop taking this drug also.

- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of new or worse liver problems like dark urine, decreased appetite, upset stomach or stomach pain, light-colored stools, any bruising or bleeding, feeling very tired, throwing up, or yellow skin or eyes.
- Shortness of breath.
- Sunburn.
- Very bad skin irritation.

- Mouth sores.
- Eye irritation.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Itching.
- Upset stomach.
- Muscle pain.
- Diarrhea.
- Headache.
- Feeling dizzy, tired, or weak.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take this drug with food.
- Swallow whole. Do not chew, open, or crush.
- Take this drug at the same time of day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.
- It is important that you do not miss or skip a dose of this drug during treatment.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- If you are not sure what to do if you miss a dose, call your doctor.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.

- Store in the original container to protect from light.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your

poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use,

available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-10-21

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Simeprevir - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center