



PATIENT & CAREGIVER EDUCATION

Spesolimab

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Spevigo

Brand Names: Canada

Spevigo

What is this drug used for?

- It is used to treat a type of psoriasis (generalized pustular psoriasis, GPP).

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have an infection.
- If you have TB (tuberculosis).

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- You will need a TB (tuberculosis) test before starting this drug.
- You may have more chance of getting an infection. Wash hands often. Stay away from people with infections, colds, or flu.
- If possible, make sure all vaccines are up to date before treatment with this drug.
- Talk with your doctor before getting any vaccines during treatment and for at least 16 weeks after your last dose of this drug. Use of some vaccines with this drug may

either raise the chance of an infection or make the vaccine not work as well.

- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.
- If you used this drug during pregnancy, tell your baby's doctor. You will need to discuss the safety and timing of certain vaccines with the doctor.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.

- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Dizziness or passing out.
- Vaginal itching or discharge.
- A severe and sometimes deadly reaction has happened. Most of the time, this reaction has signs like fever, rash, or swollen glands with problems in body organs like the liver, kidney, blood, heart, muscles and joints, or lungs. If you have questions, talk with the doctor.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling tired or weak.
- Upset stomach or throwing up.
- Diarrhea.
- Headache.
- Joint pain.
- Bruising where the shot is given.

- Swelling, warmth, or redness where this drug was given.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088.

You may also report side effects at

<https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Injection (subcutaneous):

- It is given as a shot into the fatty part of the skin on the top of the thigh or the belly area.
- If you will be giving yourself the shot, your doctor or nurse will teach you how to give the shot.
- Wash your hands before and after use.
- Do not use if the solution is cloudy, leaking, or has particles.
- This drug is clear to brownish-yellow. Do not use if the solution changes color. If you are not sure, talk with the

pharmacist.

- Do not give into skin within 2 inches (5 cm) of the belly button.
- Do not give into skin that is irritated, tender, bruised, red, scaly, hard, scarred, or has stretch marks.
- If your dose has more than 1 injection, give the injections into 2 different places at least 1 inch apart.
- Switch between the thigh and the belly area for each complete dose.
- Before using this drug, take it out of the refrigerator. Allow to reach room temperature (15 to 30 minutes). Do not heat this drug.
- Throw away after using. Do not use the device more than 1 time.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.
- If you drop the device, be sure you know if it can be used or if you need to replace it. If you are not sure, read the package insert. If you have questions, talk with the doctor or pharmacist.

Injection (I.V.):

- It is given as an infusion into a vein over a period of time.

What do I do if I miss a dose?

Injection (subcutaneous):

- Take a missed dose as soon as you think about it.
- If it is almost time for the next dose, skip the missed dose and return to your regular schedule.
- Do not take 2 doses at the same time or extra doses.
- If you are not sure what to do if you miss a dose, call your doctor.

Injection (I.V.):

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

Prefilled syringes:

- Store in a refrigerator. Do not freeze.
- Do not use if it has been frozen.
- Store in the outer carton to protect from light.

300 mg/2 mL prefilled syringes:

- If needed, you may store at room temperature up to 77°F (25°C) for up to 14 days. Write down the date you

take this drug out of the refrigerator. If stored at room temperature and not used within 14 days, throw this drug away.

Injection (I.V.):

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health

care provider.

- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any

risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-06-16

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Spesolimab - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center