

# Albuterol

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### **Brand Names: US**

ProAir Digihaler [DSC]; ProAir HFA [DSC]; ProAir RespiClick; Proventil HFA [DSC]; Ventolin HFA

### **Brand Names: Canada**

Airomir; APO-Salbutamol HFA; DOM-Salbutamol; PMS-Salbutamol; TEVA-Salbutamol; TEVA-Salbutamol HFA; Ventolin; Ventolin Diskus; Ventolin HFA; Ventolin PF [DSC]

### **What is this drug used for?**

- It is used to open the airways in lung diseases where spasm may cause breathing problems.
- It is used to prevent breathing problems that happen with exercise.

## **What do I need to tell the doctor BEFORE my child takes this drug?**

### **All products:**

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child is using another drug like this one. If you are not sure, ask your child's doctor or pharmacist.

### **All dry powder inhaler products:**

- If your child is allergic to milk, talk with the doctor.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

## **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not give more of this drug or have your child use it more often than you were told by the doctor. Deaths have happened when too much of this drug was taken. Talk with the doctor.
- Call the doctor right away if your child's normal dose does not work well, if your child's signs get worse, or if your child needs to use this drug more often than normal.
- If your child has high blood sugar (diabetes), you will need to watch your child's blood sugar closely.

### **If your child is pregnant or breast-feeding a baby:**

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

## **What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

### **All products:**

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal.
- Chest pain or pressure, a fast heartbeat, or an abnormal heartbeat.
- This drug may sometimes cause severe breathing problems. This may be life-threatening. When this

happens with an inhaler or with liquid for breathing in, most of the time it happens right after a dose and after the first use of a new canister or vial of this drug. If your child has trouble breathing, breathing that is worse, wheezing, or coughing, get medical help right away.

### **All dry powder inhaler products:**

- Pain when passing urine.
- Trouble passing urine.

### **All oral products:**

- Signs of a severe skin reaction (Stevens-Johnson syndrome, erythema multiforme) like red, swollen, blistered, or peeling skin (with or without fever); skin reaction that looks like rings; red or irritated eyes; or sores in the mouth, throat, nose, or eyes.

### **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

## **All products:**

- Feeling nervous and excitable.
- Dizziness or headache.
- Upset stomach or throwing up.
- Shakiness.

## **All products for breathing in:**

- Nose or throat irritation.
- Runny nose.

## **All dry powder inhaler products:**

- Back pain.
- Aches and pains. Mild pain drugs may help.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

## **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions

closely.

## **All inhaler products:**

- For breathing into the lungs.
- If your child is using more than 1 inhaled drug, ask your child's doctor which inhaled drug to give first.
- If giving this drug to prevent breathing problems that happen with exercise, give 15 to 30 minutes before your child exercises.
- Follow how to clean carefully.
- Put the cap back on after your child is done using a dose.

## **Inhaler (aerosol):**

- Shake the inhaler well before use.
- You will need to prime the inhaler before first use. You will also need to prime the inhaler if it has not been used for some time. Be sure you know when you need to prime the inhaler and how to do it. Talk with your doctor or pharmacist if you have questions.
- Some inhalers need to be primed if dropped. Ask your doctor or pharmacist if you are not sure.
- A spacer may be used with the inhaler for easy use.
- This inhaler may have a dose counter to keep track of

how many doses are left. If it does, throw the inhaler away when the dose counter reaches “0”.

### **All dry powder inhaler products:**

- Close the device after each dose. Do not open the device unless a dose is being used.
- Do not take the device apart or wash it. Do not use it with a spacer. Be sure your child does not breathe out into the device.
- Clean mouthpiece by wiping with a dry tissue or cloth. Do not wash or put in water.
- This inhaler has a dose counter to keep track of how many doses are left. Throw away the inhaler when the dose counter reaches “0,” 13 months after opening the foil pouch, or after the expiration date, whichever comes first.

### **Digital dry powder inhaler:**

- This product keeps track of inhaler event information and may be used with a smartphone app. This product can be used without being connected to the smartphone app. Be sure you know how to use this product and the smartphone app. Follow what the doctor has told you to do. Read the package insert for details. If you have questions, talk with the doctor.



## **Liquid for breathing in:**

- For breathing in only as a liquid (solution) by a special machine (nebulizer) into the lungs.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Do not mix other drugs in nebulizer.

## **All oral products:**

- Give this drug with or without food.

## **Oral liquid:**

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

## **What do I do if my child misses a dose?**

- If your child uses this drug on a regular basis, give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- Many times this drug is given on an as needed basis.

Do not give to your child more often than told by the doctor.

## **How do I store and/or throw out this drug?**

### **All products:**

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

### **All inhaler products:**

- Protect from cold.
- If the inhaler comes in a foil pouch, store in the foil pouch until ready for use.

### **Inhaler (aerosol):**

- Protect from heat and sunlight. Do not puncture or burn even if it seems empty.
- Throw away the inhaler after the most number of sprays have been used, even if it feels like there is

more drug in the canister.

- Store with the mouthpiece down.

### **All dry powder inhaler products:**

- Protect from heat.

### **Liquid for breathing in:**

- Store unused containers in foil pouch until use.
- Check how long vials may be stored once the pouch has been opened.

### **General drug facts**

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2024-02-15

## Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Albuterol - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center