

Carvedilol

This information from Lexicomp[®] explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Coreg; Coreg CR

Brand Names: Canada

APO-Carvedilol; Auro-Carvedilol; JAMP-Carvedilol; PMS-Carvedilol; TEVA-Carvedilol

What is this drug used for?

- It is used to treat heart failure (weak heart).
- It may be given to your child for other reasons. Talk with the doctor.
- This drug is not approved for use in children. However, the doctor may decide the benefits of taking this drug outweigh the risks. If your child has been given this drug, ask the doctor for information about the benefits and risks. Talk with the doctor if you have

questions about giving this drug to your child.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems:
Asthma, other lung or breathing problems that cause shortness of breath or wheezing, certain types of abnormal heartbeats called heart block or sick sinus syndrome, a slow heartbeat, or heart failure that is being treated with certain IV drugs.
- If your child has liver disease.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood pressure and heart rate checked often.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- To lower the chance of feeling dizzy or passing out, have your child rise slowly if your child has been sitting or lying down. Have your child be careful going up and down stairs. Dizziness and passing out may be more likely to happen when your child starts this drug or when the dose is increased.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may prevent some signs of low blood sugar like fast heartbeat. This may raise the risk of severe or long-lasting low blood sugar, especially in people with diabetes, children, and people who are fasting. This

includes people who are having surgery, are not eating like normal, or are throwing up. If you have questions, talk with the doctor.

- If your child has high blood sugar (diabetes), you will need to watch your child's blood sugar closely.
- Tell your child's doctor if your child has signs of high or low blood sugar like breath that smells like fruit, dizziness, fast breathing, fast heartbeat, feeling confused, feeling sleepy, feeling weak, flushing, headache, unusual thirst or hunger, passing urine more often, shaking, or sweating.
- Do not stop giving this drug to your child all of a sudden. If you do, chest pain that is worse and in some cases heart attack may occur. The chance may be higher if your child has certain types of heart disease. To avoid side effects, you will want to slowly stop this drug as ordered by the doctor. Call the doctor right away if your child has new or worse chest pain or if other heart problems happen.
- This drug may make it harder to tell if your child has signs of an overactive thyroid like fast heartbeat. If your child has an overactive thyroid and stops taking this drug all of a sudden, it may get worse and could be life-threatening. Talk with the doctor.
- If your child has had a very bad allergic reaction, talk

with the doctor. Your child may have a chance of an even worse reaction if your child comes into contact with what caused the allergy. If your child uses epinephrine to treat very bad allergic reactions, talk with the doctor. Epinephrine may not work as well while your child is taking this drug.

- If your child is having cataract surgery, talk with the doctor.
- If your child wears contact lenses, your child may have fewer tears or dry eyes. Call the doctor if this bothers your child.
- If your child is taking this drug and has high blood pressure, talk with the doctor before giving OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of a severe skin reaction (Stevens-Johnson syndrome) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Very bad dizziness or passing out.
- Not able to pass urine or change in how much urine is passed.
- Chest pain that is new or worse.
- Slow heartbeat.
- A burning, numbness, or tingling feeling that is not

normal.

- Feeling cold in the arms or legs.
- Change in eyesight.
- Call your child's doctor right away if your child has shortness of breath, a big weight gain, a heartbeat that is not normal, or swelling in the arms or legs that is new or worse. Worsening of heart failure has happened with this drug.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Headache.
- Diarrhea, upset stomach, or throwing up.
- Weight gain.
- Joint pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about

side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Give this drug with food.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Give this drug at the same time of day.

Extended-release capsules:

- Have your child swallow whole. Do not let your child chew, break, or crush.
- If your child cannot swallow this drug whole, you may sprinkle the contents on applesauce. If you do this, have your child swallow the mixture right away without chewing.
- If mixing on applesauce, the applesauce should not be warm. Do not sprinkle on other liquids or foods.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not put on 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not

endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-01-08

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Carvedilol - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center