

Cladribine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Mavenclad (10 Tabs); Mavenclad (4 Tabs); Mavenclad (5 Tabs); Mavenclad (6 Tabs); Mavenclad (7 Tabs); Mavenclad (8 Tabs); Mavenclad (9 Tabs)

Brand Names: Canada

Mavenclad

Warning

- This drug may lower the ability of the bone marrow to make blood cells that the body needs. If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. If you have questions, talk with the doctor.
- This drug may cause kidney problems in some patients. Talk with the doctor.
- The chance of kidney problems may be higher if your child is taking any drugs that can affect kidney function. There are many drugs that can do this. Ask the doctor or pharmacist if you are not sure.
- Very bad nerve problems have happened with this drug. This includes your child not being able to move a part of her/his body (paralysis) that can be long-lasting. Call your child's doctor if your child has any burning, numbness, tingling, or weakness of the arms or legs.

What is this drug used for?

- It is used to treat a type of leukemia.
- It may be given to your child for other reasons. Talk with the doctor.

Tablets:

- If your child has been given this form of this drug, talk with the doctor for information about the benefits and risks. Talk with the doctor if you have questions or concerns about giving this drug to your child.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Kidney disease or liver disease.
- If your child has any of these health problems: Hepatitis B, hepatitis C, HIV, or any other infection.
- If your child has TB (tuberculosis).
- If your child takes any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for hepatitis C, HIV, infections, and others. There are many drugs that must not be taken with this drug.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug. Your child may also need to avoid breast-feeding a baby for some time after the last dose. Talk with your child's doctor to see if your child needs to avoid breast-feeding a baby after the last dose.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Your child may have more chance of getting an infection. Some infections have been deadly. Have your child wash hands often. Have your child stay away from people with infections, colds, or flu.
- Your child may bleed more easily. Make sure your child is careful and avoids injury. Be sure your child has a soft toothbrush.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- If your child is not up to date with vaccines, talk with your child's doctor. Your child may need to have some vaccines before treatment with this drug.
- Talk with the doctor before your child gets any vaccines. Use of some vaccines with this drug may either raise the chance of very bad infection or make the vaccine not work as well.
- If your child has taken this drug and is getting a blood transfusion, talk with your child's doctor.
- Some products have benzyl alcohol. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol. Talk with the doctor to see if this product has benzyl alcohol in it.

If your child is or may be sexually active:

- This drug may cause harm to an unborn baby. Your child must use birth control while taking this drug. If your child gets pregnant, call your child's doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of shingles like pain, itching, or tingling; fever; muscle aches; or rash (with or without blisters).
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of a very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Feeling very tired or weak.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Fast or abnormal heartbeat.
- Patients with cancer who take this drug may be at greater risk of getting a severe health problem called tumor lysis syndrome (TLS). This may lead to death. Call your child's doctor right away if your child has a fast

or abnormal heartbeat; any passing out; trouble passing urine; muscle weakness or cramps; upset stomach, throwing up, diarrhea, or not able to eat; or feels sluggish.

- A severe brain problem called progressive multifocal leukoencephalopathy (PML) has happened with this drug. It may cause disability or can be deadly. Tell the doctor right away if your child has signs like confusion, memory problems, depression, change in the way your child acts, change in strength on 1 side is greater than the other, trouble speaking or thinking, change in balance, or change in eyesight.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Headache.
- Muscle pain.
- Irritation where this drug is given.
- Diarrhea, throwing up, upset stomach, and decreased appetite are common with this drug. If these happen, talk with your child's doctor about ways to lower these side effects. Call your child's doctor right away if any of these effects bother your child, do not get better, or are severe.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-06-24

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Cladribine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center