



PATIENT & CAREGIVER EDUCATION

About Overactive Bladder (OAB)

This information explains what overactive bladder (OAB) is, the symptoms, and how to diagnose and treat it.

About your bladder

Your bladder is an organ that holds urine (pee). As urine flows into your bladder, the walls of your bladder stretch. This helps make room for more liquid.

When you urinate, your bladder pushes urine out of your body through your urethra. (yoo-REE-thruh). The urethra is the tube that carries urine from your bladder to outside your body. While this happens, the muscles in your urethra and pelvis relax.

What is OAB?

Overactive bladder is when the muscles in your bladder squeeze down too often, too strong, or both (see figure 1). This can cause urine to come out at the wrong time. This can be uncomfortable, embarrassing, and cause you to avoid social activities.




Figure 1. Normal bladder (left) and overactive bladder (right)

Causes and risk factors of OAB

As you get older, you may be more likely to have OAB. OAB can also be caused by:

- Bladder conditions, such as infection or pelvic floor muscle dysfunction. This is when the muscles that support the bladder and other organs do not work as well as they should.
- Certain health conditions, such as diabetes.
- Diet, such as eating or drinking things that irritate (bother) your bladder. These include caffeine, spicy foods, and carbonated (fizzy) drinks.

For some people, we do not know what causes OAB.

Symptoms of OAB

People with OAB may feel the urge to pee right away or pee often. This can happen even if their bladder is not full. They may leak urine or empty their whole bladder before they can get to the bathroom.

Symptoms of OAB include:

- **Nocturia** (nok-TOOR-ee-uh): Having to pee more than once while you're sleeping.
- **Urinary urgency incontinence (UII)**: Leaking urine while having a sudden, strong urge to pee.
- **Urinary frequency**: Having to pee often, usually more than 8 times a day.
- **Urinary urgency**: Having a strong, sudden need to pee right away.

How to diagnose OAB

Talk with your healthcare provider, urogynecologist (YOO-ruh-GY-neh-KAH-loh-jist) or urologist (yoo-RAH-loh-jist) if you have symptoms of OAB.

- A urogynecologist is a doctor with special training in diagnosing and treating pelvic floor conditions, such as weak bladder.
- A urologist is a doctor with special training in diagnosing and treating urinary organs and urologic diseases. This includes pelvic floor conditions, such as weak bladder.

Your healthcare provider will do a physical exam. They will ask you when you leak urine, and how often. They may also order:

- Urodynamics (YOO-ruh-dy-NA-miks) tests to help them learn more about your bladder and urethra. A urodynamics test uses a catheter and manometer to measure the pressure in your bladder. Read *Urodynamic Evaluation* (www.mskcc.org/cancer-care/patient-education/urodynamic-evaluation) to learn more.
- Urine analysis tests to check for urinary tract infection (UTI) and blood in your urine.
- An ultrasound to see how much urine is in your bladder after you pee. Read *About Your Prostate Biopsy With Ultrasound* (www.mskcc.org/cancer-care/patient-education/about-your-prostate-

[biopsy-ultrasound](#)) to learn more.

Your healthcare provider may tell you to keep a bladder diary. They will ask you to record how much you urinate. You will record what you drink, how much, and how often. This will help them learn more about your bladder symptoms.

Treatments for OAB

Talk with your healthcare provider about which treatment or treatments for OAB are best for you.

Nutrition and diet

Some drinks can cause bladder spasms and make OAB symptoms worse. These include:

- Drinks with caffeine, such as coffee, tea, and soda.
- Drinks with fruit juice or artificial sweeteners.
- Drinks with alcohol.

Your healthcare provider can help you plan changes to your diet to ease the symptoms of OAB.

Physical therapy

Going to the bathroom often can cause your bladder to hold less urine. This can make you urinate even more often and make symptoms of OAB worse.

You can retrain your bladder by doing exercises to strengthen the muscles that help you hold in urine. Over time, this can help you go to the bathroom less often to urinate. It may take up to 3 months before you see results.

Talk with your healthcare provider to see if pelvic floor physical therapy is right for you. They may give you a referral for a pelvic floor physical therapist (PT) who can teach you these exercises.

Medicine

Your healthcare provider may prescribe medicine to help ease the symptoms of OAB. These medicines have different side effects. Talk with your healthcare provider about which medicine is right for you.

Botox

A Botox injection (shot) into your bladder wall can help to relax your bladder. This can help it hold more urine and ease the symptoms of OAB.

Your healthcare provider will give you local anesthesia (A-nes-THEE-zhuh) or sedation (seh-DAY-shun). Local anesthesia is medicine that numbs an area of your body. Sedation is when you're calm, relaxed, or sleepy from medicine you get before your injection. Then, they will use a needle with a small camera on it to inject Botox into your bladder wall.

Some people may get a UTI or have trouble emptying their bladder after getting Botox. Talk with your healthcare provider about whether Botox is right for you.

Nerve stimulation

Nerve stimulation (nerv STIM-yoo-LAY-shun) is similar to acupuncture. Your healthcare provider will put a small needle into nerves that affect bladder function. They will use a device to send small impulses into the nerves. This changes the signals your nerves send to your bladder to help you need to urinate less often. Talk with your healthcare provider about whether nerve stimulation is right for you.

Caring for your skin

For people with a vagina

All-cotton or incontinence pads, diapers, and incontinence underwear can help hold leakage. Incontinence pads are gentler on the skin than menstrual pads for people with a vagina.

Keep the skin around the vagina dry whenever possible. It may help to use a barrier cream, such as Aquaphor® or Vaseline®, can help.

For people with a penis

All-cotton or incontinence pads, diapers, and incontinence underwear can help hold leakage. Penile clamps may be helpful for people with a penis.

When to call your healthcare provider

If you think you have symptoms of a UTI, call your healthcare provider right away. This includes:

- Feeling burning when you urinate.
- Having urine that smells very bad.
- Having cloudy urine.

You should also call your healthcare provider if you have:

- A fever of 100.4 °F (38 °C) or higher.
- Blood in your urine when you pee.
- Trouble emptying your bladder.
- Other symptoms of overactive bladder.
- Body shaking.
- Chills.
- Confusion, if you are 80 years old or older.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

About Overactive Bladder (OAB) - Last updated on February 13, 2024

All rights owned and reserved by Memorial Sloan Kettering Cancer Center