



PATIENT & CAREGIVER EDUCATION

Side Effects of Palliative Radiation Therapy to the Abdomen or Pelvis

Memorial Sloan Kettering nurse practitioner Carol Ann Milazzo describes side effects commonly associated with radiation therapy to the abdomen or pelvis. Radiation can cause patients to feel temporarily fatigued. Because radiation travels through the skin, it may cause the treated area to look and feel sunburned – a condition called radiation dermatitis.

Patients may want to ask their doctors about how to manage these potential side effects during and immediately after radiation. Other common side effects may include nausea, heartburn, diarrhea, and a sensation of burning hemorrhoids.



Please visit www.mskcc.org/videos/side-effects-palliative-radiation-therapy-abdomen-pelvis to watch this video.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Side Effects of Palliative Radiation Therapy to the Abdomen or Pelvis - Last updated on September 1, 2013

All rights owned and reserved by Memorial Sloan Kettering Cancer Center